

Extracellular Lipase of Malassezia as Anti Dandruff Drug Target: A Review

by Wong Hendra Wijaya

Submission date: 07-Sep-2022 03:21PM (UTC+0700)

Submission ID: 1894263684

File name: 196-1601307335.pdf (781.64K)

Word count: 5078

Character count: 28959

Extracellular Lipase of *Malassezia* as Anti Dandruff Drug Target: A Review

Wong H. Wijaya¹, Kris H Timotius^{2*} and Jonathan K. Wijaya¹

¹Department of Dermatology, Faculty of Medicine, Krida Wacana Christian University (UKRIDA), Jakarta, Indonesia

²Department of Biochemistry, Faculty of Medicine, Krida Wacana Christian University (UKRIDA), Jakarta, Indonesia

Corresponding author: Kris H Timotius

Email: kh_timotius@ukrida.ac.id

ABSTRACT

This review seeks to answer how is the adaptation of *Malassezia* spp. for their indwelling life in scalp; what are the roles of extracellular enzymes as a virulence factor in the pathogenesis of *Malassezia* for dandruff/seborrheic dermatitis; and how can Indonesian herbal materials be screened as potential anti-dandruff drug by observing their ability to inhibit extracellular lipase of *Malassezia*. Three species of *Malassezia* (*M. furfur*, *M. globosa* and *M. restricta*) are commonly indwelling in the scalp-skin either in normal healthy individuals, or in patients with dandruff/seborrheic dermatitis. The infectious lipolytic *Malassezia* spp. are adapted with the scalp-skin environment and depend significantly on fatty acid supplement. To be able to live in the scalp, extracellular lipases are produced by *Malassezia*. Their invasive pathogenicity is dependent on high activities and production of extracellular lipases which plays a significant role in the survival and virulence of *Malassezia*. Therefore, these lipases may become the main target in screening for anti-dandruff drugs. For this purpose, several Indonesian herbal materials can be screened due to their potentials as lipase inhibitor.

Keywords: Dandruff Lipase, Herbal material, *Malassezia*, Seborrheic dermatitis

Correspondence:

Kris H Timotius

Department of Biochemistry, Faculty of Medicine, Krida Wacana Christian University (UKRIDA), Jakarta, Indonesia

Email: kh_timotius@ukrida.ac.id

Dandruff causes discomfort in scalp and creates an undesired social image. Two factors are important for the aetiology of Dandruff and Seborrheic Dermatitis (D/SD): physiological factors (sebum, pH, water content, and individual susceptibility) and microbial metabolism (especially *Malassezia* spp.). The lipophilic yeast of the skin microbiome, *Malassezia* spp., determine the severity of the D/SD. (1, 2). Several species of *Malassezia* are involved in skin/scalp disorders including D/SD, which together affect >50% of humans (3)

In this review, three research questions are discussed. The first research question is how the predominant species of *Malassezia* adapt to the D/SD scalp and whether there are differences between *Malassezia* as natural flora of the skin and as D/SD causing pathogen. The second research question is about the extracellular enzymes as a virulence factor of *Malassezia*. *Malassezia* species, even though acts as a natural inhabitant of the healthy skin, can cause several skin diseases. Its infection and colony formation are a result of complex interactions between the *Malassezia* and its extracellular enzymes as its virulence factor (4). The third research question is whether extracellular lipases of *Malassezia* can become target of anti-dandruff drug. The interaction between these enzyme and their inhibitor(s) is important to be understood (5-7) because this alternative approach can lead to the discovery of potential novel anti-dandruff drugs in the future. So far, the treatment of dandruff relies mainly on azole anti-fungal drugs, which are nowadays characterized by emerging resistance, poor compliance, clinical inefficacy and several side effects. It is necessary to promote the alternative for potential novel anti-dandruff drugs (8-10). Several Indonesian herbal materials have potential in anti-lipase activity and therefore can be used as anti-dandruff drug in the future.

The aims of this study were to find characteristics of the scalp predominant indwelling *Malassezia* spp. which cause D/SD, and to describe their extracellular enzymes as a

virulence factor which determines the survival and pathogenesis of *Malassezia*; and to select Indonesian herbal materials which have potential as anti-lipase activity and are prospective to become anti-dandruff drugs. After the introduction, the following sections are presented: the adaptation of *Malassezia* to the scalp environment (D/SD); the extracellular enzymes as a virulence factor of *Malassezia*; and Indonesian herbal materials as a source of lipase inhibitors. At the end, are sections for conclusions and recommendations.

THE ADAPTATION OF MALASSEZIA TO THE SCALP ENVIRONMENT

Scalp environment¹

The scalp is characterized with rich sebum production and carbohydrate deficient environment. The pH of the scalp plays an important physiological role. The pH of the stratum corneum is acidic, between 4.1 to 5.8. This pH is determined by several mechanisms/factors, such as the amount of fatty acids produced, products of filaggrin degradation, activated sodium-hydrogen exchanger, and released melanosome. The acidic pH of the stratum corneum of the scalp inhibits the formation of microbial colony (e.g. by *Staphylococcus aureus* and *Malassezia*). However, the acidic pH also influences skin condition, lipid biosynthesis, cells aggregation, epidermal differentiation and/or desquamation. The disturbance of skin barrier by the increased pH can cause skin inflammation and other epidermal diseases. Topical treatment to normalize the pH helps to maintain a healthy physiological microbiota, as well as the condition of the skin barrier, the induction of epidermal differentiation and avoiding skin inflammation (11).

Lipid-dependence of *Malassezia*

Human pathogenic *Malassezia* spp. is lipid-dependent and are adapted to the narrow niche on sebum-rich skin. Most of the *Malassezia* showed an absolute requirement for

long fatty acid chain. In the case of the three predominant dandruff-causing species of *M. furfur*, *M. globosa* and *M. restricta*, oleic acid alone can initiate dandruff-like desquamation (1, 12). The lipid dependence of *Malassezia* can be understood because of the absence of fatty acid synthase gene. Their incapability in fatty acids biosynthesis are complemented by the presence of multiple secreted lipases to aid in degrading host lipid and harvesting the resulted fatty acids. Therefore, high levels of lipase is a probable importance in their pathogenicity, and expressed on human scalp (1, 13). *Malassezia* has many genes encoding the secretion of hydrolases (e.g., acid sphingomyelinase, aspartyl protease, lipase, phospholipase) which are important indicators of host-specific adaptation (3). Therefore, it can be concluded that the presence and activity of multiple secreted lipases are important to harvest the host's lipids (14).

The genomic analysis shows that a larger set of genes are lost. This evidence is concordant with the adaptation of *Malassezia* to skin's carbohydrate-deficient scalp environment. Therefore, it is important for *Malassezia* to produce and secrete extracellularly for the several key enzymes, such as aspartyl protease, lipases, phospholipases, and other peptidases. A recent genomic resource explained the development of *Malassezia* niche-specificity and potential virulence, as well as the abundance and distribution in the environment and on human skin (15).

The long-chain fatty acids are not synthesized by *Malassezia*. Therefore, the biosynthesis of myristic acid requires the addition of preformed fatty acids as its precursor. The growth media of *Malassezia* needs to be supported by the addition fatty acids with a carbon chain length of greater than 10. The additional supply, including the fatty acids are odd- or even-numbered carbon chain lengths. The kind of added lipid that is used during growth, affects the fatty acid composition of the *Malassezia*. This means that the fatty acids are used as energy sources and also incorporated directly into cellular lipids. The lipids present on normal human scalps are able to fulfill the lipid requirement of the *Malassezia* (3).

Inability of *Malassezia* to ferment glucose/carbohydrate

Malassezia is unable to ferment sugars as the sole source of carbon but are able to use lipid. Beside sebum production, scalp environment is characterized by carbohydrate deficient environment. Carbohydrates, electrolytes, vitamins and trace elements are not required for the growth of *Malassezia*. *Malassezia* is normally able to grow in vitro under aerobic, microaerophilic as well as anaerobic conditions (11).

THE EXTRACELLULAR LIPASE OF MALASSEZIA AS MAIN INVASIVE VIRULENCE FACTORS AND AS TARGET OF NEW ANTI-DANDRUFF DRUG

Main invasive virulence factors

The pathogenesis of *Malassezia* depends on several virulence factors, namely its ability to produce reactive oxygen species (ROS) (16), allergens (17), metabolites (indole and azelaic acid) (18, 19), unique volatile gamma lactones, and extracellular enzymes (Lipase, Carbonic anhydrase, and Protease). The survival and growth of *Malassezia* in the scalp, as well as subsequent scalp damage, are mediated by its virulence factors which set them apart from harmless microbiota. The host-fungus

interactions lead to the development of D/SD (20). The pathogenic role of *Malassezia* are determined by host immune system and the virulence factors. Several virulence factors in *Malassezia* are their enzymatic lipolytic activity that result in the production of distinct metabolites, special cell wall features, and irritant metabolic by products including bioactive indole derivatives (21, 22). Compared with lipase, carbonic anhydrase and protease are less important factors to be used as target of the anti-dandruff drug. The first attention should be given to lipase which enables *Malassezia* to use triglyceride sebum in the scalp. The secretion of enzymes by *Malassezia* is considered as an important factor in the invasion and dissemination in the host. Therefore, it is suggested that secreted lipases are responsible for the pathogenicity of *Malassezia* spp. However, the expression of these genes is not yet investigated during the development of the disease. The *M. globosa* lipase gene expression on human scalp is still in the early step. (3). The role of lipase activity is very crucial in the pathogenicity of *M. globosa* towards dandruff and seborrheic dermatitis (D/SD).

Lipases are produced by *Malassezia* with the help of lipoygenase and are used to digest triglyceride of the sebum. These enzymes are located on the cell wall and the cytoplasmic membrane. The greatest lipase activity is observed during logarithmic phase where the cells are active growing. The activity is influenced by the concentration of the substrate. (23) At least three different extracellular lipases are known as essential for the growth of *Malassezia*. In vitro growth of *Malassezia* species also produce phospholipase. Activity of phospholipase releases arachidonic acid that are involved in inflammation of the skin. The activity of phospholipase is the basic mechanism by which *Malassezia* species may trigger inflammation. Isolates of *M. furfur*, *M. globosa*, and *M. restricta*, also produce esterase, phosphatase-acid and Naphtol-AS-BI-phosphohydrolase in significant amounts from most isolates excepted for *M. restricta*. *M. restricta* has limited enzymatic activity compared to other species. Five extracellular enzymes including keratinase (protease), lipase and phospholipase are produced (24, 25). Genetic analysis of the *M. restricta* genome identified at least eight lipase sequences. The lipases MrLip1, MrLip2 and MrLip3 are tested for their activity using mono-, di- and triacylglycerol substrates. Hydrolysed by the *M. restricta* lipase MrLip1 and MrLip2 are limited to the mono- and diacylglycerol. Meanwhile, MrLip3 hydrolyses all three substrates. Lipases are responsible for the hydrolysis of triacylglycerols, the main component of human sebum. The information regarding lipases from *M. restricta* may aid in the search for anti-dandruff agents (26). Even though most lipases of *Malassezia* are considered to be active against triacylglycerides, there is a small group of lipases that are active only on mono- and diacylglycerides (27) (Table 1).

Lipase of *M. furfur*

The activity of lipase can be measured in the extract of the soluble and insoluble cells, as well as in the supernatant of the culture. The enzymatic activity and cell growth are first induced and later inhibited by increasing concentrations of Tween-80 (polyethylene-sorbitan monooleate). The characters of the lipolytic system explained the incremental growth of *M. furfur* that alter the surface lipid in the scalp, such as seborrheic dermatitis. The lipolytic system is induced by the substrate and influenced by the environmental pH and/or different cations). (23). The

growth of *M. furfur* is totally dependent on the external lipid supply. With the lipolytic activity, Malassezia hydrolyses lipid and release fatty acids that are necessary for both growth and pathogenicity. The sequence similarity of the gene MFLIP1 is high compared with the other known extracellular lipases (28). The hyphal form of Malassezia shows higher lipase activities than those of the yeast form (29).

Lipase of *M. globosa*

Malassezia globosa is characterised by its highest lipase activity compared to the rest of the Malassezia species. It has 6 phospholipase-C encoding genes which secrete lipases, and two phospholipase-D genes and one phospholipase-B gene which do not secrete. (3). MgLIP1 and MgLIP2 are found in *M. globosa*. The activity of lipase from *M. globosa* (SMG1) is special because of its specific activity on hydrolysis mono- and diacylglycerol, but not on triacylglycerol (29). *M. globosa* lipase (SMG1) has an unusual activity in which it is capable in catalysing epoxidation of alkenes. The long chain carboxylic acids are catalysed to peroxides that subsequently react with double bonds of alkenes to produce epoxides. SMG1 is selectively reactive to carboxylic acids. Since the activity of SMG1 lipase is regulated by allosteric mechanism upon binding to the lipophilic-hydrophilic phase interface, it is used to drive the epoxidation in the lipophilic phase exclusively (30). SMG1 is one of the few members in the fungal lipase family that strictly specific for mono- and diacylglycerol (31). The diglyceride-hydrolyzing lipase of *M. globosa* can be isolated. It is able to degrade diolein, but not triolein. But the lipase of the intact cells show activity against both substrates. This evidence indicates that the presence of at least another lipase is necessary. The lipase gene expression can be detected on the human scalp by reverse transcription-PCR. This step is important in molecular description of lipid metabolism on the scalp, ultimately leading toward a test of its role in D/SD aetiology (32).

Lipase of *M. restricta*

The presence of lipases and phospholipases have significant contribution in the virulence of *M. restricta* as being the most frequently isolated Malassezia spp. from the human skin. These lipases in the host scalp environment are significantly able to produce free fatty acids for Malassezia (33). Therefore, extracellular lipases of Malassezia play a critical role in its survival on the host skin surface.

Targeting Malassezia lipase for discovering new anti-dandruff drugs

Two approaches are known in discovering anti-dandruff drugs which have inhibitory activity on the extracellular lipase of Malassezia, namely with docking analysis at screening bioactivity compounds from herbal materials. In order to improve the eradication rate of Malassezia, the use of extracellular enzymes, such as lipase and /or carbonic anhydrase (MgCA), is carried out. This topic is important in dealing with the latest discoveries of herbal medicine for the management of Malassezia spp.-related lipases. New opportunities are open for discovering innovative and alternative anti-dandruff drugs in the future (8). In this review, we focus on herbal materials, mainly from Indonesia, which have anti-lipase activity since lipase is the most important target. The resistance to single and/or multi-drug is greatly hampering patient management. Additionally, drug

resistance mechanisms include altered drug-target interactions, reduced cellular drug concentrations mediated by drug efflux transporters, and permeability barriers associated with the presence of biofilms. Therefore, the development of better diagnostic tools and approaches that allow targeted use of antifungals are essential to enhance drug efficacy (28)

Currently, the most utilized anti-dandruff drug is the Azole group, such as ketoconazole and climbazole. Ketoconazole significantly inhibits fungal filamentation or mycelia formation. Since mycelia typify the pathogenic form of Malassezia infection, the capacity of ketoconazole to block morphogenesis may represent an additional important effect of the antifungal (29). Climbazole is an example of an imidazole antifungal agent that can provide anti-dandruff benefits when incorporated into a hair shampoo. Azole antifungals are able to prevent the ergosterol synthesis (30). However, anti-fungal therapies, as in the case of Malassezia related dandruff, are considered less effective due to of chronic character of Malassezia infections and high percentage of relapses. Therefore, there is a urgent need to develop new approach for Malassezia therapies, in this case, with novel inhibitors of lipase of Malassezia (31).

THE INDONESIAN HERBAL MATERIALS WITH ANTI-LIPASE POTENTIAL

Natural remedies for treating dandruff are becoming more popular. Natural antimicrobials/antifungal are importance in recent times to combat the global challenge of antibiotic resistance. Herbal antimicrobial materials are gaining importance in therapy because their ability to encounter minimal challenges of emergence of resistance. As example, embelin of *Embelia ribes* was identified as an anti-lipase and fungi static agent. Embelin exhibits a synergistic effect with the anti-fungal drug ketoconazole (KTZ) against Malassezia spp. (9). Several Indonesian herbal materials that have the potential as anti-dandruff agents are listed in table 2.

- *Aloe vera* is known for hair growth, acne, wound healing, and shampoo. It is able to inhibit adipose lipase activities. And therefore, *Aloe vera* has great potential as functional foods in the activation adipose lipolysis and the prevention of obesity-related metabolic alterations. *Aloe vera* gel has inhibitory potential against pancreatic lipase activity, and significant inhibition on pancreatic lipase (32, 33).
- *Azadirachta indica* (neem tree) contains azadirachtin as its main constituent in various parts of the tree. Azadirachtin, a tetranortriterpenoid, is remarkable for its chemical complexity and for its biological activity. It is reported that azadirachtin inhibit the rice bran lipase, an esterase, which is a versatile enzyme that catalyzes the hydrolysis of ester linkages, primarily in neutral lipids such as triglycerides (34, 35)
- *Centella asiatica* is used to as a remedy for acne because of its ability to inhibit lipase produced by the bacteria responsible for acne formation. It is reported that pancreatic lipase inhibitory activity of *C. asiatica* extract was significantly higher than rutin but lower than orlistat, an anti-obesity drug (36).
- *Nigella sativa* is known for its anti-fungal capacity. Ethanol seed extract of *Nigella sativa* has anti-fungal capacity against *M. furfur*. It was identified that *N. sativa* contains bioactive constituents which it

stimulates further studies on the dermatological effects and application of *N. sativa*(37).

- *Musa paradisiaca* is popular for its fruit everywhere in the world. Anti-fungal and antibiotic principles are found in the peel and pulp of fully ripe bananas. It can be used effectively for a beauty treatment either for skin or hair. Bananas have the ability to improve manageability and shine while moisturizing our skin and helping to prevent dandruff. Therefore, it is good to use it as an excellent treatment for in addressing the dandruff's problem and in retaining moisture on our scalp(38).
- *Eleusine indica* showed pancreatic lipase inhibitory activity with no significant difference between its methanol extract and the standard drug Orlistat. It is a good component of anti-dandruff shampoo(39).
- *Ruta angustifolia* is reported as herbal material for various diseases, including dandruff. The antibacterial and antifungal effects of four traditional plants essential oils(40)

CONCLUSIONS AND FUTURE DIRECTIONS (RECOMMENDATIONS)

Conclusions

- The lipolytic *Malassezia* is adapted to the scalp environment, in order to transform their existence as skin flora normal to a pathogenic agent. They have lost their lipid biosynthetic genes and are significantly dependent on fatty acid supplement.
- As one of the virulence factors of *Malassezia*, extracellular lipase is important for its survival in the scalp environment. Therefore, key extracellular lipases of *Malassezia* can become target of dandruff drugs.
- Indonesian herbal materials are potential and promising as a source of novel discovered anti-dandruff drugs.

Recommendation for future direction

- Further research on interaction host-microbe in the scalp environment is recommended.
- Lipase as virulence factors of *Malassezia* should be understood properly in order to support the effort to search novel anti-*Malassezia*, especially anti-dandruff drugs. The additional information on lipase formation in needed to aid the screening of anti-dandruff agents.
- Extracellular lipase produced by *Malassezia* is a good target of seeking novel anti-dandruff drug. Searching and screening natural bioactive compounds can be started by exploring lipase inhibitors. Investigation about lipase inhibitor on fungal virulence factors is necessary in order to identify the precise mechanism of antifungal activity and to do clinical trials on *Malassezia*

20

Conflict of Interest : The authors declared no conflict of interest.

Ethical clearance : No ethical clearance was needed for this literature review.

Source of funding : No special funding was obtained for this literature review.

REFERENCES

1. Dawson TL, Jr. *Malassezia globosa* and *restricta*: breakthrough understanding of the etiology and treatment of dandruff and seborrheic dermatitis through whole-genome analysis. The journal of investigative dermatology Symposium proceedings. 2007;12(2):15-9.
2. Meray Y, Gencalp D, Guran M. Putting It All Together to Understand the Role of *Malassezia* spp. in Dandruff Etiology. *Mycopathologia*. 2018.
3. Xu J, Saunders, C.W., Hu, P., Grant, R. A., Boekhout, T., Kuramae, E. E. et al. . Dandruff-associated *Malassezia* genomes reveal convergent and divergent virulence traits shared with plant and human fungal pathogens. *Proceedings of the National Academy of Sciences of the United States of America*. 2007;104(47):18730-5.
4. Angiolella L, Leone C, Rojas F, Mussin J, de Los Angeles Sosa M, Giusiano G. Biofilm, adherence, and hydrophobicity as virulence factors in *Malassezia furfur*. *Medical mycology*. 2018;56(1):110-6.
5. Capasso C, Supuran CT. Bacterial, fungal and protozoan carbonic anhydrases as drug targets. *Expert opinion on therapeutic targets*. 2015;19(12):1689-704.
6. Del Prete SV, D. Osman, S. M. AlOthman, Z. Capasso, C. Supuran, C. T. Anion inhibition studies of the dandruff-producing fungus *Malassezia globosa* beta-carbonic anhydrase MgCA. *Bioorganic & medicinal chemistry letters*. 2015;25(22):5194-8.
7. Entezari Heravi YB, S. Nocentini, A. Del Prete, S. Saboury, A. A. Sereшти, H. et al , Capasso C, Gratteri P, Supuran CT. Inhibition of *Malassezia globosa* carbonic anhydrase with phenols. *Bioorganic & medicinal chemistry*. 2017;25(9):2577-82.
8. Angiolella LC, S., Maccallini, C., Giusiano, G. Supuran, C. T. Targeting *Malassezia* species for Novel Synthetic and Natural Antidandruff Agents. *Current medicinal chemistry*. 2017;24(22):2392-412.
9. Sivasankar CG, S. Bhaskar, J. P. Krishnan, V. Pandian, S. K. Evaluation of selected Indian medicinal plants for antagonistic potential against *Malassezia* spp. and the synergistic effect of embelin in combination with ketoconazole. *Microbial pathogenesis*. 2017;110:66-72.
10. Supuran CT. Bortezomib inhibits bacterial and fungal beta-carbonic anhydrases. *Bioorganic & medicinal chemistry*. 2016;24(18):4406-9.
11. Proksch E. pH in nature, humans and skin. *The Journal of dermatology*. 2018.
12. Theelen BC, C. Gaitanis, G. Bassukas, I. D. Boekhout, T. Dawson, T. L., Jr. *Malassezia* ecology, pathophysiology, and treatment. *Medical mycology*. 2018;56(suppl_1):10-25.
13. Aspiroz CA, M. Varea, M. Rezusta, A. Rubio, C. . Isolation of *Malassezia globosa* and *M. sympodialis* from patients with pityriasis versicolor in Spain. *Mycopathologia*. 2002;154(3):111-7.
14. Guillot J, Hadina S, Gueho E. The genus *Malassezia*: old facts and new concepts. *Parassitologia*. 2008;50(1-2):77-9.
15. Deng Y, Liu X, Wu J, Lee J, Chen S, Cheng Y, et al. The host plant metabolite glucose is the precursor of diffusible signal factor (DSF) family signals in *Xanthomonas campestris*. *Applied and environmental microbiology*. 2015;81(8):2861-8.

16. Spater S, Hipler UC, Haustein UF, Nenoff P. [Generation of reactive oxygen species in vitro by *Malassezia* yeasts]. *Der Hautarzt; Zeitschrift für Dermatologie, Venerologie, und verwandte Gebiete*. 2009;60(2):122-7.
17. Gioti AN, B. Li, W. Xu, J. Andersson, A. Averette, A. F., Munch K, Wang X, Kappauf C, Kingsbury JM, Kraak B, et al. Genomic insights into the atopic eczema-associated skin commensal yeast *Malassezia sympodialis*. *mBio*. 2013;4(1):e00572-12.
18. Gaitanis GM, P. Stathopoulou, K. Bassukas, I. D. Alexopoulos, E. C., Velegriaki A, Skaltsounis AL. AhR ligands, malassezin, and indolo[3,2-b]carbazole are selectively produced by *Malassezia furfur* strains isolated from seborrheic dermatitis. *The Journal of investigative dermatology*. 2008;128(7):1620-5.
19. Ashbee HR A-AZ, Z. Whitehead Z, Evans E G V. Proc Fourth Congr Eur Confed Med Mycol, abstr 5, . 1998.
20. Brunke S, Mogavero S, Kasper L, Hube B. Virulence factors in fungal pathogens of man. *Current opinion in microbiology*. 2016;32:89-95.
21. Cafarchia C, Otranto D. The pathogenesis of *Malassezia* yeasts. *Parassitologia*. 2008;50(1-2):65-7.
22. Hort W, Mayser P. *Malassezia* virulence determinants. *Current opinion in infectious diseases*. 2011;24(2):100-5.
23. Plotkin LIS, L. Mathov, I. Galimberti, R. Leoni, J. . Characterization of the lipase activity of *Malassezia furfur*. *Journal of medical and veterinary mycology : bi-monthly publication of the International Society for Human and Animal Mycology*. 1996;34(1):43-8.
24. Mancianti FR, A. Nardoni, S. Corazza, M. Extracellular enzymatic activity of *Malassezia* spp. isolates. *Mycopathologia*. 2001;149(3):131-5.
25. Muhsin TM, Aubaid AH, al-Duboon AH. Extracellular enzyme activities of dermatophytes and yeast isolates on solid media. *Mycoses*. 1997;40(11-12):465-9.
26. Sommer B, Overy DP, Kerr RG. Identification and characterization of lipases from *Malassezia restricta*, a causative agent of dandruff. *FEMS yeast research*. 2015;15(7).
27. Lan D, Wang Q, Xu J, Zhou P, Yang B, Wang Y. Residue Asn277 affects the stability and substrate specificity of the SMG1 lipase from *Malassezia globosa*. *International journal of molecular sciences*. 2015;16(4):7273-88.
28. Perlín DS, Rautemaa-Richardson R, Alastruey-Izquierdo A. The global problem of antifungal resistance: prevalence, mechanisms, and management. *The Lancet Infectious Diseases*. 2017;17(12):e383-e92.
29. Youngchim SN, J. D. Chongkae, S. Vanittanokom, N. Ketoconazole inhibits *Malassezia furfur* morphogenesis in vitro under filamentation optimized conditions. *Archives of dermatological research*. 2017;309(1):47-53.
30. Perez-Rivera AAH, T. Aardema, M. J. Nash, J. F. Evaluation of the genotoxicity of the imidazole antifungal climbazole: comparison to published results for other azole compounds. *Mutation research*. 2009;672(1):27-39.
31. Guo SH, W. Zhang, J. Wang, Y. Novel inhibitor against *Malassezia globosa* LIP1 (SMG1), a potential anti-dandruff target. *Bioorganic & medicinal chemistry letters*. 2015;25(17):3464-7.
32. Rahoui W, Merzouk H, El Hacı IA, Bettioui R, Azzi R, Benali M. Beneficial effects of Aloe vera gel on lipid profile, lipase activities and oxidant/antioxidant status in obese rats. *Journal of Functional Foods*. 2018;48:525-32.
33. Taukooerah U, Mahomoodally MF. Crude Aloe vera Gel Shows Antioxidant Propensities and Inhibits Pancreatic Lipase and Glucose Movement In Vitro. *Advances in pharmacological sciences*. 2016;2016:3720850.
34. Gangadhara, Ramesh Kumar P, Prakash V. Inhibition of rice bran lipase by azadirachtin from *Azadirachta indica*. *Journal of the Science of Food and Agriculture*. 2009;89(10):1642-7.
35. Gangadhara PRK, Prakash V. Inhibition of rice bran lipase by azadirachtin from *Azadirachta indica*. *J Sci Food Agric*. 2009;89:1642-7.
36. Supkamonseni N, Thinkratok A, Meksuriyen D, Srisawat R. Hypolipidemic and hypoglycemic effects of *Centella asiatica* (L.) extract in vitro and in vivo. *Indian journal of experimental biology*. 2014;52(10):965-71.
37. Aljabre SHM, M. Alakloby O, A. Randhawa M. Dermatological affects of *Nigella sativa*. *Journal of Dermatology & Dermatologic Surgery*. 2015;19.
38. Kaminaro SS. Pengaruh minyak buah pisang (*Musa paradisiaca* L.) terhadap pengurangan ketombe pada kulit kepala. *Jurnal Tata Rias Universitas Negeri Jakarta*. 2015;8 (8) 31-5.
39. Lai YS, Y. Lee, al. JKe. In vitro lipase inhibitory effect of thirty two selected plants in Malaysia. *Evidence-Based Complementary and Alternative Medicine*. 2014.
40. Haddouchi F, Chaouche TM, Zaouali Y, Ksouri R, Attou A, Benmansour A. Chemical composition and antimicrobial activity of the essential oils from four *Ruta* species growing in Algeria. *Food chemistry*. 2013;141(1):253-8.
41. Brunke S, & B. Hube. MfLIP1, a gene encoding an extracellular lipase of the lipid-dependent fungus *Malassezia furfur*. *Microbiology (Reading, England)*. 2006;152(Pt 2):547-54.
42. Juntachai W, Kajiwarara S. Differential Expression of Extracellular Lipase and Protease Activities of Mycelial and Yeast Forms in *Malassezia furfur*. *Mycopathologia*. 2015;180(3-4):143-51.
43. DeAngelis YS, CW Johnstone, KR Reeder, NL Coleman, CG, Kaczvinsky Jr JR, Gale C, Walter R, Mekel M, Lacey MP, Keough TW, et al. Isolation and expression of a *Malassezia globosa* lipase gene, LIP1. *Journal of Investigative Dermatology*. 2007;127:2138-46.
44. Juntachai W, Oura T, Kajiwarara S. Purification and characterization of a secretory lipolytic enzyme, MgLIP2, from *Malassezia globosa*. *Microbiology (Reading, England)*. 2011;157(Pt 12):3492-9.
45. Sommer BO, D. P. Haltli, B. Kerr, R. G. Secreted lipases from *Malassezia globosa*: recombinant expression and determination of their substrate specificities. *Microbiology (Reading, England)*. 2016;162(7):1069-79.
46. Park MC, Y. J. Lee, Y. W. Jung, W. H. Whole genome sequencing analysis of the cutaneous pathogenic yeast *Malassezia restricta* and identification of the major lipase expressed on the scalp of patients with dandruff. *Mycoses*. 2017;60(3):188-97.
47. Naga PP, K. A, K. D. Comparison of potency of antifungal action of dandruff shampoos and different plant extracts. *International Journal of Medical Research & Health Sciences*. 2015;4(2):327-31.

48. Mahashu Vellingiri DTPaJMS. Effects of processing conditions on the stability of polyphenolic contents and antioxidant capacity of *Dolichos lablab* L. . Journal of food science and technology. 2013;50(4):731-8.
49. Sahak MKA KN, Abbas G, Drasman S, Hashim NH, Adil DSH. The role of *Nigella sativa* and its active constituents in learning and memory. Evidence-Based Complementary and Alternative Medicine. 2016;2016 6 pages.
50. Perera A, Ton SH, Palanisamy UD. Perspectives on geraniin, a multifunctional natural bioactive compound. Trends in Food Science & Technology. 2015;44(2):243-57.

Table 1. List of lipases produced by *Malassezia*

Producer	Type of lipase	Substrate	pH	Ref.
<i>M.furfur</i>	MfLIP1	MAGs, TAGs	Opt. 4.5-5.8 Range 3.5-7.0 Inactive >6.5	(41) (42)
<i>M.globosa</i>	MgLIP1	MAGs, DAGs	Opt. 5.5	(43)
	MgLIP2	MAGs, DAGs	No at 8	(44)
	MgLIP3	MAGs, DAGs, TAGs		(45)
<i>M.restricta</i>	MrLip1	MAGs, DAGs	Opt. 5,5	(26)
	MrLip2	MAGs, DAGs		(46)
	MrLip3	MAGs, DAGs, TAGs		
	MrLip4			
	MrLIP5			

Table 2. List of Indonesian herbal materials which are potential candidate for lipase inhibitor

Name of plant	Local name	Part used	Ref.
<i>Aloe vera</i> ,	<i>Lidah buaya</i>	Leaf	(47)
<i>Azadirachta indica</i> ,	<i>Neem, nimba</i>	Leaf	(35,47)
<i>Centella asiatica</i>	<i>Goto kola, Pegagan</i>	Leaf	(36)
<i>Coriandum sativum</i>	<i>Coriander, Ketumbar</i>	Leaf	(47)
<i>Dolichos lablab</i>	<i>Daun komak</i>	Leaf	(48)
<i>Eleusine indica</i>	<i>Goose grass, Rumput belulang</i>		(39)
<i>Mentha asiatica</i>	<i>Mint</i>	Leaf	(47)
<i>Musa paradisiaca</i>	<i>Pisang</i>	Fruit, peel	(38)
<i>Nigella sativa</i>	<i>Biji jintan hitam</i>	Seed	(37) (49)
<i>Ocimum tenuiflorum</i>	<i>Tulsi</i>	Leaf	(47)
<i>Piper bettle</i>	<i>Betel leaf</i>	Leaf	(47)
<i>Punica granatum</i>	<i>Delima</i>	Leaf	(50)
<i>Ruta angustifolia</i>	<i>Daun inggu</i>	Leaf	(40)

Extracellular Lipase of Malassezia as Anti Dandruff Drug Target: A Review

ORIGINALITY REPORT

20%
SIMILARITY INDEX

17%
INTERNET SOURCES

19%
PUBLICATIONS

4%
STUDENT PAPERS

PRIMARY SOURCES

1 cmr.asm.org Internet Source **2%**

2 bjp.sagepub.unboundmedicine.com Internet Source **2%**

3 repository.unair.ac.id Internet Source **2%**

4 www.idexlab.com Internet Source **2%**

5 lib.bioinfo.pl Internet Source **2%**

6 pubmed.ncbi.nlm.nih.gov Internet Source **1%**

7 Submitted to Institute of International Studies Student Paper **1%**

8 Www.Fulltxt.Org Internet Source **1%**

9 insights.ovid.com Internet Source **1%**

10 Letizia Angiolella, Simone Carradori, Cristina Maccallini, Gustavo Giusiano, Claudiu T. Supuran. " Targeting species for Novel Synthetic and Natural Antidandruff Agents ", Current Medicinal Chemistry, 2017
Publication 1 %

11 www.labome.org
Internet Source 1 %

12 Wiebke Hort. "Malassezia virulence determinants :", Current Opinion in Infectious Diseases, 04/2011
Publication 1 %

13 synapse.koreamed.org
Internet Source 1 %

14 Changjin Ji, Na Liu, Jie Tu, Zhuang Li, Guiyan Han, Jian Li, Chunquan Sheng. "Drug Repurposing of Haloperidol: Discovery of New Benzocyclane Derivatives as Potent Antifungal Agents against Cryptococcosis and Candidiasis", ACS Infectious Diseases, 2019
Publication 1 %

15 Clio Dessinioti, Andreas Katsambas. "Seborrheic dermatitis: Etiology, risk factors, and treatments:", Clinics in Dermatology, 2013
Publication 1 %

16 Tingting Xu, Lu Liu, Shulin Hou, Jinxin Xu, Bo Yang, Yonghua Wang, Jinsong Liu. "Crystal structure of a mono- and diacylglycerol lipase from *Malassezia globosa* reveals a novel lid conformation and insights into the substrate specificity", *Journal of Structural Biology*, 2012
Publication 1 %

17 worldwidescience.org
Internet Source 1 %

18 www.chemistry.uoc.gr
Internet Source 1 %

19 Chandran Sivasankar, Subramanian Gayathri, James Prabhanand Bhaskar, Venkat Krishnan, Shunmugiah Karutha Pandian. "Evaluation of selected Indian medicinal plants for antagonistic potential against *Malassezia* spp. and the synergistic effect of embelin in combination with ketoconazole", *Microbial Pathogenesis*, 2017
Publication 1 %

20 www.nature.com
Internet Source 1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On