

# Irritable bowel

*by* Ina S Timan

---

**Submission date:** 26-Oct-2022 09:02AM (UTC+0700)

**Submission ID:** 1935521590

**File name:** Irritable\_bowel.pdf (482.75K)

**Word count:** 5171

**Character count:** 24086



Jornal de  
**Pediatria**

www.jpmed.com.br



## ORIGINAL ARTICLE

## 1 Irritable bowel syndrome in Indonesian adolescents ☆

3 Q3 Yudianita Kesuma <sup>a</sup>, Rini Sekartini <sup>b</sup>, Ina S. Timan <sup>c</sup>, Agnes Kurniawan <sup>d,e</sup>,  
4 Saptawati Bardosono <sup>e</sup>, Agus Firmansyah <sup>b</sup>, Yvan Vandenplas <sup>f,\*</sup>

5 Q4 <sup>a</sup> Sriwijaya University and Moh Hoesin Hospital, Faculty of Medicine, Department of Pediatrics, Palembang, Indonesia

6 <sup>b</sup> University of Indonesia and Cipto Mangunkusumo Hospital, Faculty of Medicine, Department of Pediatrics, Jakarta, Indonesia

7 <sup>c</sup> University of Indonesia Cipto Mangunkusumo Hospital, Faculty of Medicine, Department of Clinical Pathology, Jakarta, Indonesia

8 <sup>d</sup> University of Indonesia, Faculty of Medicine, Department of Parasitology, Jakarta, Indonesia

9 <sup>e</sup> University of Indonesia, Faculty of Medicine, Department of Nutrition, Jakarta, Indonesia

10 <sup>f</sup> Free University of Brussels, UZ Brussel Hospital, KidZ Health Castle, Brussels, Belgium

11 Received 7 October 2019; accepted 6 February 2020

## KEYWORDS

12 Intestinal integrity;  
13 Irritable bowel  
14 syndrome;  
15 Quality of life;  
16 Adolescent

## 1 Abstract

2 **Objective:** Irritable bowel syndrome is a frequent functional gastrointestinal disorder. The aims  
3 of this study were to investigate its epidemiology, focusing on the role of intestinal mucosal  
4 integrity and to evaluate the impact on the quality of life.

5 **Methods:** A community-based survey applying a comparative cross sectional approach was  
6 conducted in six high schools in Palembang. Subjects were recruited using multistage random  
7 sampling divided in two groups. Rome III criteria were used to establish a diagnosis of IBS  
8 in combination with a questionnaire to determine risk factors. Determination of fecal alpha-  
9 1-antitrypsin and calprotectin <sup>16</sup> levels was performed to determine impaired intestinal mucosal  
10 integrity. A questionnaire <sup>16</sup> was used to evaluate how quality of life was affected by irritable  
11 bowel syndrome.

12 **Results:** The survey was performed in 454 14–18 years old adolescents, of whom 30.2% fulfilled  
13 the Rome III criteria for IBS, with the following subtypes: 36.5% diarrhea, 18.9% constipa-  
14 tion, 21.9% mixed, and 22.6% unclassified. Major risk factors were female gender, bullying,  
15 age 14–16 years, history of constipation and diarrhea, eating nuts, and drinking coffee, tea,  
16 and soft drinks. There was a significant association with intestinal inflammation ( $p = 0.013$ ). A  
17 significantly <sup>32</sup> impaired quality of life was found ( $p = 0.001$ ).

18 **Conclusions:** The prevalence of irritable bowel syndrome in adolescents was high, with bullying,  
19 female gender, age 14–16 years, constipation and diarrhea, and dietary consumption of soft  
20 drinks, coffee, and tea as risk factors. A significant association with intestinal inflammation was  
21 found.

22 © 2020 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open  
23 access article under the CC BY-NC-ND license ([http://creativecommons.org/licenses/by-nc-nd/  
24 4.0/](http://creativecommons.org/licenses/by-nc-nd/4.0/)).

25 ☆ Please cite this article as: Kesuma Y, Rini Sekartini R, Timan I.S., Kurniawan A, Bardosono S, Firmansyah A et al. Irritable bowel syndrome  
26 in Indonesian adolescents. J Pediatr (Rio J). 2020. <https://doi.org/10.1016/j.jpmed.2020.02.001>

27 Corresponding author.

28 E-mail: [yvan.vandenplas@uzbrussel.be](mailto:yvan.vandenplas@uzbrussel.be) (Y. Vandenplas).

29 <https://doi.org/10.1016/j.jpmed.2020.02.001>

30 0021-7557/© 2020 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND  
31 license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Adolescence is a period during which various important changes happen in a child's journey towards adulthood.<sup>1-4</sup> Irritable bowel syndrome (IBS) is a common functional gastrointestinal disorder. It is characterized by abdominal pain that usually improves with defecation. The onset is associated with a change in stool consistency or frequency that cannot be explained by structural or biochemical abnormalities. IBS is characterized by several subtypes: diarrhea, constipation, mixed, or unclassified.<sup>5-7</sup> The pathophysiology of IBS still remains unknown. Data from other studies found structural changes such as abnormal permeability and intestinal inflammation in IBS patients.<sup>8,9</sup> Several case control studies reported that risk factors of IBS are related to age, gender, socio-economic status, education level, 'Western lifestyle,' increased personal hygiene, population overgrowth, stress, ethnicity, climate, medications, past history of dysentery,<sup>20</sup> sexual abuse, and other psychosocial factors.<sup>5,10-13</sup> IBS is a major public health problem, with an important impact on health care cost, activity limitation, quality of life (QoL), and behavior disorders.<sup>14-17</sup>

The present study aimed to characterize IBS in adolescents regarding prevalence, risk factors, and subtypes, to analyze the involvement of intestinal mucosal integrity, and to determine the impact of IBS on QoL.

## Methods

A community-based survey with comparative cross-sectional design was conducted in 14-18 year old high school students. Subjects were recruited using multistage random sampling from six high schools from six districts in Palembang, between April and September of 2017. The study consisted of two stages: the first stage was a cross-sectional survey, and the second stage, a cross-sectional comparison. Regarding the first stage, it was estimated that this study would require 355 subjects, according to the estimated prevalence of IBS of 21%<sup>12</sup>, with a permissible error of 5%. The first stage was intended to determine the prevalence, risk factors, and subtypes of IBS. Regarding the second part of the study, participants were randomly classified into an IBS and a non-IBS group. The number was based on a sample size calculated with a formula for comparing two populations, in order to analyze the association of IBS with gut integrity and the quality of life. This calculation resulted in the finding that 140 adolescents, divided into 70 IBS and 70 non-IBS from the first part of the study, was the number needed in order to analyze the association of IBS with gut integrity and QoL (Fig. 1).

The inclusion criteria were adolescents in presumed good health. Subjects with any chronic disease such as asthma, diabetes mellitus, or cardiac, renal, or hematological




Fig. 1 Recruitment of subjects.

IBS, irritable bowel syndrome; QoL, quality of life; AAT, alpha-1-antitrypsin; CP, calprotectin.

## Irritable bowel syndrome in adolescents

diseases were excluded. Those who had received antibiotics or antiparasitic treatments (metronidazole, tinidazole, albendazole) in the last two weeks were also excluded. All subjects who met the inclusion criteria were included in the study after obtaining the written or oral consent of a parent or guardian. Information was obtained from all participants regarding identity and presence of symptoms, allowing the presence of Rome III criteria to be determined, and symptoms related to QoL were elucidated by direct interview. The QoL assessment was performed using the IBS-QoL questionnaire to evaluate the impact of IBS. The IBS-QoL questionnaire was validated in a preliminary study in 30 adolescents, prior to this study.

All participants also underwent routine physical examination and anthropometric measurements. Nutritional status was defined according to the Centers for Disease Control and Prevention (CDC) 2000 growth charts: obese if body mass index (BMI) > P95; overweight if BMI > P85; underweight if BMI for age < P5; severe underweight if BMI for age < P3.

The diagnosis of IBS was established according to the Rome III criteria and was confirmed by a pediatric gastroenterologist using the same criteria. Participants were classified into an IBS and a non-IBS group, in order to analyze the association of IBS with gut integrity. Fecal ELISA DRG<sup>®</sup> was used to measure alpha-1-antitrypsin (AAT) and fecal calprotectin (CP) to determine gut integrity. Data were analyzed using SPSS v. 22.0. Normally distributed data are shown as the mean  $\pm$  standard deviation. Student's *t*-test for independent samples, the Mann-Whitney test, Pearson's chi-squared test, and Fisher's exact test were used whenever appropriate. Analysis for CP and AAT levels are presented as median, with minimum and maximum range. Since the distribution of the CP and AAT levels was not normal according to Kolmogorov-Smirnov, the Mann-Whitney test was used. A *p*-value  $\leq$  0.05 was considered statistically significant (95% confidence interval).

This research was approved by the Health Research Ethics Committee, Faculty of Medicine, University of Indonesia, under protocol No. 774/UN2.F1/ETIK/2017, of August 21st 2017.

## Results

From April to September 2017, this study enrolled 454 high school students, representing 37.2% boys and 62.8% girls, with an average age of  $15.8 \pm 0.97$  years. Most students (70.3%) had good nutritional status; the rest were malnourished. IBS according to Rome III criteria was found in 137 students (30.2%), with the following distribution by subtype: diarrhea IBS (IBS-D) subtype was the most frequent, followed by unclassified IBS (IBS-U), mixed IBS (IBS-M), and constipation IBS (IBS-C), with percentages of 36.5%; 22.6%; 21.9%; 18.9%, respectively (Table 1).

## Risk factors for IBS

Risk factors related to IBS were identified (Table 1), including female gender ( $p=0.001$ , OR 2.13, 95% CI 1.369–3.326), grumpy father ( $p=0.040$ , OR 1.71, 95% CI 1.053–2.774), sexual abuse ( $p=0.008$ , OR 3.38, 95% CI 2.934–3.902), bullying ( $p=0.034$ , OR 2.67, 95% CI 1.148–6.212), eating spicy

soup ( $p=0.043$ , OR 1.56, 95% CI 1.034–2.339), eating nuts ( $p=0.002$ , OR 1.97, 95% CI 1.309–2.954), consumption of soft drinks ( $p=0.003$ , OR 1.95, 95% CI 1.271–2.988), and history of diarrhea ( $p=0.020$ , OR 1.83, 95% CI 1.124–2.987) and constipation ( $p=0.023$ , OR 2.03, 95% CI 1.139–3.626).

## Association between IBS and gut integrity

According to sample size calculation, 70 patients were needed in each group. Gut integrity was estimated by measuring intestinal permeability and inflammation using stool assays. Intestinal permeability was represented by fecal AAT concentration and inflammation by fecal CP. The cut-off point for AAT was  $> 43.7$  mg/dL and for CP it was  $> 50$  mg/kg. The study results showed that fecal AAT levels were not normally distributed; data are shown as median (min-max). AAT concentration in the IBS group was decreased compared with the non-IBS group (57.44 [0.0–541.50] vs. 66.88 [1.24–525.30]), but there was no significant relationship between fecal AAT and IBS ( $p=0.361$ ). However, fecal CP was increased in the IBS group (120.77 [0.70–3,255.0] vs. 88.62 [6.88–3,842.5]) and there was a significant relationship between fecal CP and IBS ( $p=0.013$ ; Table 2).

## The impact IBS on QoL

The IBS-Quality of Life questionnaire (IBS-QoL) was developed by Patrick et al.<sup>16</sup> This study had shown that, after a standard translation and linguistic validation and reliability questionnaire IBS linguistic validation method, the Indonesian IBS-QoL can be administered in other populations with appropriate reliability and validity.

Permission was obtained from the original questionnaire developer and the MAPI Research Trust (Lyon, France) to develop this study.<sup>16</sup> The present authors followed the recommended process of the MAPI Research Trust, as well as the principles ordered by the translation standard forward-backward translation was done; two health professionals did the forward translation independently and backward translation was done by two professional translators separately as well.

The Indonesian IBS-QoL was validated by 30 subjects with a simple random method.<sup>17</sup> Reliability analyses demonstrated high internal consistency and reproducibility of the Indonesian IBS-QoL for the overall score, as well as for subscale scores, comparable to the original study (Cronbach's alpha = 0.95, ICC = 0.88).

The QoL of subjects in IBS group was lower than the non-IBS group. IBS-QoL among the IBS group showed lower QoL than the non-IBS group ( $n=140$ ). The negative impact of IBS on QoL was shown in this study. There was a significant relationship with low quality of life ( $p=0.001$ ; mean difference = 5.8; 95% CI = 2.816–12.083; Table 3).

## Discussion

This study aimed to determine the epidemiology of IBS and to identify relationships of gut integrity and QoL in adolescents with IBS. IBS is reported to be one of the most common functional gastrointestinal disorders in children.

Q1 Table 1 Assessment of the risk factors related to irritable bowel syndrome (IBS) (n = 104).

Characteristic	IBS (+) n (%)	IBS (-) n (%)	Total	p-Value	OR	95% CI
Age						
14–16 Y	110 (80.3)	225 (70.9)	335	0.051 <sup>a</sup>	1.67	1.025 – 2.708
17–18 Y	27 (19.7)	92 (29.1)	119			
Gender						
Girls	102 (74.5)	183 (57.7)	285	0.001 <sup>a</sup>	2.13	1.369 – 3.326
Boys	35 (25.2)	134 (42.5)	169			
Nutritional status						
Malnourished	37 (27.0)	98 (30.9)	135	0.469 <sup>a</sup>	0.83	0.529 – 1.292
Good	100 (73.0)	219 (69.1)	319			
Gestational age						
Preterm	3 (2.2)	5 (1.6)	8	0.702 <sup>b</sup>	1.40	0.329 – 5.929
Term	134 (97.8)	312 (98.4)	446			
Birth weight						
< 2500 g	2 (1.5)	3 (0.9)	5			
> 2500 g	135 (98.5)	314 (99.1)	449	0.640 <sup>b</sup>	1.56	0.256 – 9.386
No. of siblings						
<2	35 (25.5)	79 (24.9)	114			
>2	102 (74.5)	238 (75.3)	340	0.981 <sup>a</sup>	1.03	0.652 – 1.639
Birth order						
<3	109 (79.6)	277 (87.4)	386			
>3	28 (20.4)	40 (12.6)	68	0.046 <sup>a</sup>	0.56	0.330 – 0.956
Marital status						
Married	116 (84.7)	283 (89.3)	399			
Divorced	21 (15.3)	34 (10.7)	55	0.221 <sup>*</sup>	0.66	0.370 – 1.192
Mother's behavior						
Grumpy	45 (32.8)	83 (26.2)	128			
Patient	92 (67.2)	234 (73.8)	326	0.182 <sup>a</sup>	1.38	0.892 – 2.132
Father's behavior						
Grumpy	35 (25.5)	53 (16.7)	88			
Patient	102 (74.5)	264 (83.3)	366	0.040 <sup>a</sup>	1.71	1.053 – 2.774
Psychosocial problems						
- Having been assaulted	24 (37.5)	40 (62.5)	64	0.219 <sup>b</sup>	1.47	0.847 – 2.553
- Sexual abuse	4 (100)	0 (0)	4	0.008 <sup>b</sup>	3.38	2.934 – 3.902
- Fighting	29 (31.2)	64 (68.8)	93	0.912 <sup>b</sup>	1.06	0.648 – 1.738
- Bullying	12 (52.2)	11 (47.8)	23	0.034 <sup>a</sup>	2.67	1.148 – 6.212
- Trauma	18 (30.0)	42 (70.0)	60	1.000 <sup>a</sup>	0.99	0.548 – 1.791
- Negative emotions	33 (31.4)	72 (68.6)	105	0.843 <sup>a</sup>	1.08	0.674 – 1.730
- Abuse	2 (100)	0 (0)	2	0.091 <sup>b</sup>	3.35	2.907 – 3.856
Diet						
- Fish cakes	101 (31.0)	225 (69.0)	326	0.629 <sup>a</sup>	1.15	0.731 – 1.801
- Sweet layer cakes	6 (31.6)	13 (68.4)	19	1.000 <sup>a</sup>	1.07	0.398 – 2.879
- Spicy soup (pindang)	84 (34.4)	160 (65.6)	244	0.043 <sup>a</sup>	1.56	1.034 – 2.339
- Steamed spicy fish	22 (29.7)	52 (70.3)	74	1.000 <sup>a</sup>	0.98	0.566 – 1.680
- Milk	100 (31.1)	222 (68.9)	322	0.599 <sup>a</sup>	1.16	0.740 – 1.809
- Canned foods	72 (31.7)	155 (68.3)	227	0.540 <sup>a</sup>	1.16	0.775 – 1.729
- Processed meats	60 (31.1)	133 (68.9)	193	0.794 <sup>a</sup>	1.08	0.719 – 1.616
- Nuts	70 (38.9)	110 (61.1)	180	0.002 <sup>a</sup>	1.97	1.309 – 2.954
- Fried Snacks	125 (31.3)	274 (68.7)	399	0.226 <sup>a</sup>	1.60	0.813 – 3.138
- Tea	81 (31.0)	180 (69.0)	261	0.719 <sup>a</sup>	1.10	0.733 – 1.653
- Coffee	29 (26.6)	80 (73.4)	109	0.417 <sup>a</sup>	0.79	0.491 – 1.288
- Soft drinks	96 (35.7)	173 (64.3)	269	0.003 <sup>a</sup>	1.95	1.271 – 2.988
Eating duration						
< 10 minutes	10 (7.3)	24 (7.6)	34	1.000	0.96	0.447 – 2.069

Table 1 (Continued)

Characteristic	IBS (+) n (%)	IBS (-) n (%)	Total	p-Value	OR	95% CI
> 10 minutes	127 (92.7)	293 (92.4)	420			
Previous illness						
Diarrhea	35 (41.2)	50 (58.8)	85	0.020 <sup>a</sup>	1.83	1.124 – 2.987
Constipation	24 (44.4)	30 (55.6)	54	0.023 <sup>a</sup>	2.03	1.139 – 3.626
Food allergies	22 (34.4)	50 (58.8)	72	0.520 <sup>a</sup>	1.25	0.716 – 2.193
Family history illness						
Recurrent abdominal pain	6 (7.4)	4 (3.57)	10	0.073 <sup>b</sup>	3.58	0.995 – 12.909
Subtype of IBS						
IBS Diarrhea	50 (36.5)					
IBS Constipation	26 (18.9)					
IBS Mixed	30 (21.9)					
IBS Unclassified	31 (22.6)					

Chi-squared test (a: continuity correction, b: Fisher's exact test).

Table 2 Association of gut integrity with irritable bowel syndrome (IBS; n = 140).

	IBS (+) n = 70	IBS (-) n = 70	p
AAT (n/%)	38 (48.7)	40 (51.3)	0.361
Median (min-max)	57.44 (0.0–541.5)	66.88 (1.2–525.3)	
CP (n/%)	60 (85.7)	50 (71.4)	
Median (min-max)	120.77 (0.70–3255.0)	88.62 (6.68–3842.0)	0.013

AAT, alpha-1-antitripsin; CP, calprotectin; statistics: Mann-Whitney test.

Table 3 Association of irritable bowel syndrome (IBS) with quality of life (QoL).

	IBS (+) n = 70	IBS (-) n = 70	p	Mean difference	95 % CI
Total IBS QoL	92.34 ± 12.6	97.95 ± 5.9	<0.001	5.6	2.3–8.9
Mean + SD					
Dysphoria	94.46 ± 13.5	96.96 ± 7.1	0.029	2.5	1.1–6.1
Mean + SD					
Interference with activity	91.07 ± 14.4	97.70 ± 6.3	< 0.001	6.6	2.9–10.4
Mean + SD					
Sexual	97.86 ± 8.2	99.64 ± 2.9	0.001	1.8	0.3–3.9
Mean + SD					
Health worry	89.88 ± 15.9	97.62 ± 7.1	< 0.001	7.7	3.6–11.9
Mean + SD					
Food avoidance	88.09 ± 19.9	97.85 ± 7.6	< 0.001	9.8	4.7–14.8
Mean + SD					
Social reaction	92.05 ± 15.4	98.12 ± 6.9	< 0.001	6.1	2.1–10.1
Mean + SD					
Relationships	92.74 ± 14.1	97.74 ± 8.9	0.001	5.0	1.0–8.9
Mean + SD					
Body image	92.59 ± 13.8	97.95 ± 7.7	0.001	5.4	2.3–8.9
Mean + SD					

Student's *t*-test, unpaired.

The prevalence of IBS in this study was higher than reported in studies from North America, Iran, Korea, Japan, and Singapore, where the prevalence varied between 8% and 22%.<sup>10–13,18,19</sup> Meta-analyses reported an overall prevalence of 8.8%, 12.4% in Asia, 13% in China, 23% in Turkey, and 2.8–5.1% in the United States.<sup>5,7,10–12,18</sup> It is unclear whether

these differences are due to differences in interpretation of Rome criteria due to cultural differences or if they represent real differences. In the United States, Self et al. reported that IBS-C was present in 58.1%, IBS-D in 5.4%, and IBS-M in 36.5%.<sup>19</sup> The result of the present study was similar to the data from Rajindrajith & Devanarayana in Sri Lanka, who

also reported a predominant distribution of IBS-D.<sup>6</sup> In contrast, a prospective hospital-based study from Italy noted IBS-C as the most prevalent IBS subtype (45%). According to an Asian study, an almost similar prevalence was noted for the different subtypes: IBS-C, IBS-D, and IBS-M each occurred in 29–30%.<sup>7,10</sup> The prevalence of subtypes of IBS is likely to depend on the methodology, but also on regional differences.

IBS is more frequent in 14–16-year-old adolescents, than in 16–18-year-olds, and more common in girls (62.8%) than boys. Gender difference is not found in every study on IBS, but if a gender difference is reported, it is always more prevalent in girls, with odds ratios between 1.2 and 3.1<sup>5–7,13,20</sup> in Western countries and 0.8–2.1 in developing countries.<sup>20</sup> Studies revealed that IBS is more common in females than in males, with a ratio of 2–2.5:1. Gender related cultural issues can play a role in this. Differences in access to health care and cultural factors such as help-seeking behavior may contribute to the prevalence differences as well.<sup>7,10–13,21</sup>

According to the logistic regression analysis, the main risk factors were bullying, gender, age, previous history of constipation or diarrhea, nut consumption, and consumption of coffee, tea, juice, and soft drinks, with odds ratios of 1.81–2.86. The present study did not find socio-economic status and education level of the parents to be risk factors for IBS. Multistage and multiphase random sampling were implemented.

This study measured intestinal permeability (fecal AAT) and inflammation (fecal CP) by ELISA. Fecal AAT level had no significant relation with IBS, which differs from the findings by Lisowska-Myjak et al., who reported a relation between fecal AAT levels and IBS diarrhea.<sup>9</sup> AAT is a marker of intestinal mucosal damage.<sup>22</sup> Fecal CP was related to IBS ( $p=0.013$ ). Choi & Jeong<sup>23</sup> reported an elevated stool CP in 35% of adolescents with IBS ( $p<0.01$ ). Choi & Jeong also reported that the mean concentration of fecal CP was significantly higher in children and adolescents with IBS than in healthy children ( $83.50 \pm 164.70$  vs.  $17.77 \pm 10.92$ ;  $p<0.001$ ). Several studies use fecal CP as a tool to differentiate organic disease from functional disorders.<sup>23,24</sup> This finding may be related to environmental enteric dysfunction, which is caused by poverty, poor hygiene and sanitation, climate, infection, as well as vitamin A and zinc deficiency.<sup>25,26</sup>

As expected, the QoL of adolescents with IBS was lower than in those without IBS ( $92.34 \pm 12.6$  vs.  $97.95 \pm 5.9$ ;  $p=0.001$ ). Studies from France,<sup>15</sup> the United States, and Korea all confirm these findings.<sup>27,28</sup> IBS is known to interfere with the physical aspects of health-related QoL, including daily activities and work productivity, yet limited data is available on the effect of the IBS subtypes on daily activity.

There are several strengths and novelties in the present research. Firstly, this research is a large scale community study regarding IBS, prevalence of the subtypes, and relation to gut integrity. It is known that IBS has a negative impact on the QoL of adolescents. Secondly, this study proved the possibility of an organic etiology of IBS symptoms in Indonesian adolescents, which substantiates findings from developing tropical countries to European and American definitions, proposing Rome III criteria as the basis for the diagnosis of

IBS, defined as a functional recurrent abdominal pain condition.

The limitation in this study was its relatively small sample size to analyze gut integrity. Further studies with larger sample sizes and long-term follow-up are needed. Moreover, the Rome III criteria were applied, while the updated Rome IV criteria have been developed. Additionally, the severity of IBS was not related to the impact on QoL.

## Conclusion

The prevalence of IBS, predominantly the IBS-D subtype, was higher compared with other studies, as well in developed and developing countries. The prevalence of IBS in adolescents in Palembang was as high as 30.2%. This study proved the existence of organic involvement (inflammation of the intestinal mucosa). More studies are needed to further explore these findings.

## Conflicts of interest

The authors declare no conflicts of interest.

## References

- WHO Expert Committee on Child Mental Health and Psychosocial Development, World Health Organization. <https://apps.who.int/iris/handle/10665/41266>, 1977.
- Indonesia. Undang-Undang Republik Indonesia Nomor 23 Tahun 2002 Tentang Perlindungan Anak. Republik Indonesia: Kementerian Pemberdayaan Perempuan; 2002.
- Feigelman S, Marcel VA. Growth, development and behavior. In: Nelson Kliegman, editor. Text Book of Pediatric. 18th ed. Philadelphia: Saunders/Elsevier publication; 2007. p. 145–78.
- Soeroso S. Masalah Kesehatan Remaja. Sari Pediatri. 2016;3:189–97.
- Devanarayana NM, Rajindrajith S. Irritable bowel syndrome in children: current knowledge, challenges and opportunities. World J Gastroenterol. 2018;24:2211–35.
- Rajindrajith S, Devanarayana NM. Subtypes and symptomatology of irritable bowel syndrome in children and adolescents: a school-based survey using Rome III criteria. J Neurogastroenterol Motil. 2012;18:298–304.
- Devanarayana NM, Rajindrajith S, Pathmeswaran A, Abegunasekara C, Gunawardena NK, Benninga MA. Epidemiology of irritable bowel syndrome in children and adolescents in Asia. J Pediatr Gastroenterol Nutr. 2015;60:792–8.
- Shulman RJ, Eakin MN, Czyzewski DI, Jarrett M, Ou CN. Increased gastrointestinal permeability and gut inflammation in children with functional abdominal pain and irritable bowel syndrome. J Pediatr. 2008;153:646–50.
- Lisowska-Myjak B, Jacek M, Zborowska H. Search for the laboratory parameters of inflammation in the serum and faeces of patients with irritable bowel syndrome. Gastroenterol Res Pract. 2016;2016:1–11, 730986.
- Chang FY, Lu CL, Chen TS. The current prevalence of irritable bowel syndrome in Asia. J Neurogastroenterol Motil. 2010;16:389–400.
- Chey WD, Kurlander J, Eswaran S. Irritable bowel syndrome: a clinical review. JAMA. 2015;313:949–58.
- Kumar A, Ashwlayan VD, Verma M. Irritable bowel syndrome: a review. Acta Scient Pharma Sciences. 2018;2:104–14.

- 326 13. Zhu X, Chen W, Zhu X, Shen Y. A cross-sectional study of risk  
327 factors for irritable bowel syndrome in children 8–13 years of  
328 age in Suzhou, China. *Gastroenterol Res Pract.* 2014;2014:1–6.
- 329 14. Adeniyi OF, Adenike Lesi O, Olatona FA, Esezobor CI, Ikobah JM.  
330 Irritable bowel syndrome in adolescents in Lagos. *Pan Afr Med*  
331 *J.* 2017;28:93.
- 332 15. Lee EH, Kwon O, Hahm KB, Kim WH, Kim JI, Cheung DJ, et al.  
333 Irritable bowel syndrome-specific health-related quality of life  
334 instrument: development and psychometric evaluation. *Health*  
335 *Qual Life Outcomes.* 2016;14:22.
- 336 16. Patrick DL, Drossman DA, Frederick IO, Dicesare J, Puder  
337 KL. Quality of life in persons with irritable bowel syndrome:  
338 development and validation of a new measure. *Dig Dis Sci.*  
339 1998;43:400–11.
- 340 17. Bjorner JB, Baum RR, Carson RT, Boer R, Tucker C, Andrae  
341 DA, et al. Mapping the irritable bowel syndrome quality of life  
342 measure (IBS-QOL) to EQ-5D in patients with irritable bowel  
343 syndrome with diarrhoea (IBS-D). *Value in Health.* 2016;19:  
344 A379.
- 345 18. Hyams JS, Di Lorenzo C, Saps M, Shulman RJ, Staiano A,  
346 van Tilburg M. Childhood functional gastrointestinal disorders:  
347 child/adolescent. *Gastroenterology.* 2016;150:1456–68.
- 348 19. Self MM, Czyzewski DI, Chumpitazi BP, Weidler EM, Shulman RJ.  
349 Subtypes of irritable bowel syndrome in children and adoles-  
cents. *Clin Gastroenterol Hepatol.* 2014;12:1468–73.
20. Kim YS, Kim N. Sex-gender differences in irritable bowel syn- 350  
drome. *J Neurogastroenterol Motil.* 2018;24:544–58. 351
21. Fukudo S, Kaneko H, Akiho H, Inamori M, Endo Y, Okumura T, 352  
et al. Evidence-based clinical practice guidelines for irritable 353  
bowel syndrome. *J Gastroenterol.* 2015;50:11–30. 354
22. Lussier B, Wilson AA. Alpha-1 antitrypsin: the protein. In: 355  
Wanner A, Sandhaus RA, editors. *Alpha-1 Antitrypsin.* Cham: 356  
Springer International Publishing; 2016. p. 17–30. 357
23. Choi YJ, Jeong SJ. Is fecal calprotectin always normal in children 358  
with irritable bowel syndrome? *Intest Res.* 2019;17:546. 359
24. Melchior C, Aziz M, Aubry T, Gourcerol G, Quillard M, Zalar 360  
A, et al. Does calprotectin level identify a subgroup among 361  
patients suffering from irritable bowel syndrome? Results of a 362  
prospective study. *United Europ Gastroenterol J.* 2017;5:261–9. 363
25. Harper KM, Mutasa M, Prendergast AJ, Humphrey J, Manges AR. 364  
Environmental enteric dysfunction pathways and child stunting: 365  
a systematic review. *PLoS Negl Trop Dis.* 2018;12:e0006205. 366
26. Crane RJ, Jones KD, Berkley JA. Environmental enteric dysfunction: 367  
an overview. *Food Nutr Bull.* 2015;36:576–87. 368
27. Singh P. Patients with irritable bowel syndrome-diarrhea have 369  
lower disease-specific quality of life than irritable bowel 370  
syndrome-constipation. *World J Gastroenterol.* 2015;21:8103. 371
28. Park JM, Choi M-G, Kim YS, Choi CH, Choi SC, Hong SJ, et al. 372  
Quality of life of patients with irritable bowel syndrome in 373  
Korea. *Qual Life Res.* 2009;18:435–46. 374

UNCORRECTED PROOF



# Irritable bowel

---

## ORIGINALITY REPORT

---

**20%**  
SIMILARITY INDEX

**13%**  
INTERNET SOURCES

**13%**  
PUBLICATIONS

**6%**  
STUDENT PAPERS

---

## PRIMARY SOURCES

---

**1** [www.pubfacts.com](http://www.pubfacts.com) **2%**  
Internet Source

---

**2** Yudianita Kesuma, Agus Firmansyah, Saptawati Bardosono, Ika Puspa Sari, Agnes Kurniawan. "Blastocystis ST-1 is associated with Irritable Bowel Syndrome-diarrhoea (IBS-D) in Indonesian adolescences", Parasite Epidemiology and Control, 2019 **2%**  
Publication

---

**3** Submitted to Isfahan University of Medical Sciences **2%**  
Student Paper

---

**4** Submitted to Adtalem Global Education **1%**  
Student Paper

---

**5** [bsdwebstorage.blob.core.windows.net](http://bsdwebstorage.blob.core.windows.net) **1%**  
Internet Source

---

**6** [ecc.isc.gov.ir](http://ecc.isc.gov.ir) **1%**  
Internet Source

---

**7** [www.hindawi.com](http://www.hindawi.com) **1%**  
Internet Source

---

8	Young Sun Kim, Nayoung Kim. "Sex-Gender Differences in Irritable Bowel Syndrome", Journal of Neurogastroenterology and Motility, 2018 Publication	1 %
9	Submitted to Western Governors University Student Paper	1 %
10	coek.info Internet Source	1 %
11	dspace.uib.es Internet Source	1 %
12	academic.oup.com Internet Source	1 %
13	www.wjgnet.com Internet Source	1 %
14	Jae Myung Park. "Quality of life of patients with irritable bowel syndrome in Korea", Quality of Life Research, 05/2009 Publication	<1 %
15	You Jin Choi, Su Jin Jeong. "Is fecal calprotectin always normal in children with irritable bowel syndrome?", Intestinal Research, 2019 Publication	<1 %
16	www.science.gov Internet Source	<1 %

17	Internet Source	<1 %
18	Submitted to University of Western Sydney Student Paper	<1 %
19	Niranga Manjuri Devanarayana, Shaman Rajindrajith. "Irritable bowel syndrome in children: Current knowledge, challenges and opportunities", World Journal of Gastroenterology, 2018 Publication	<1 %
20	"Poster Session Clinical", European Journal of Heart Failure, 2013. Publication	<1 %
21	"50th ESPGHAN Annual Meeting", Journal of Pediatric Gastroenterology & Nutrition, 2017 Publication	<1 %
22	<a href="http://www.nature.com">www.nature.com</a> Internet Source	<1 %
23	<a href="http://www.pubmedcentral.nih.gov">www.pubmedcentral.nih.gov</a> Internet Source	<1 %
24	Submitted to University of Alabama Student Paper	<1 %
25	<a href="http://www.acarindex.com">www.acarindex.com</a> Internet Source	<1 %

26 Xueping Zhu, Weichang Chen, Xiaoli Zhu, Yueping Shen. "A Cross-Sectional Study of Risk Factors for Irritable Bowel Syndrome in Children 8–13 Years of Age in Suzhou, China", *Gastroenterology Research and Practice*, 2014  
Publication <1 %

---

27 [issuu.com](http://issuu.com)  
Internet Source <1 %

---

28 [trialsjournal.biomedcentral.com](http://trialsjournal.biomedcentral.com)  
Internet Source <1 %

---

29 [www.cureus.com](http://www.cureus.com)  
Internet Source <1 %

---

30 [www.irjournal.org](http://www.irjournal.org)  
Internet Source <1 %

---

31 [www.mdpi.com](http://www.mdpi.com)  
Internet Source <1 %

---

32 [dokumen.pub](http://dokumen.pub)  
Internet Source <1 %

---

33 [www.omicsonline.org](http://www.omicsonline.org)  
Internet Source <1 %

---

34 [kanazawa-u.repo.nii.ac.jp](http://kanazawa-u.repo.nii.ac.jp)  
Internet Source <1 %

---

Exclude quotes On

Exclude matches Off

Exclude bibliography On