

B javanica

by Adelina Simamora

Submission date: 17-May-2020 08:17PM (UTC+0700)

Submission ID: 1326201359

File name: PharmacognJ-11-3-479_B_Javanica_Published.pdf (707.83K)

Word count: 5980

Character count: 32061

Antidiabetic, Antibacterial and Antioxidant Activities of Different Extracts from *Brucea javanica* (L.) Merr Seeds

Adelina Simamora^{1,*}, Kris Herawan Timotius¹, Adit Widodo Santoso²

ABSTRACT

Introduction: The seeds of *B. javanica* are known as herbal material for anticancer, antimalaria and antiamoeba. Limited report is available on their antidiabetic and antibacterial properties. **Methods:** Methanol (ME) and Ethyl acetate extracts (EAE) were studied for their inhibition activities on α -glucosidase *in vitro*, including combination of ME with acarbose and inhibition mechanism. Antibacterial activity was tested by well diffusion and microdilution methods. The extracts were evaluated for their *in vitro* antioxidant property by DPPH assay, as well as their phenolic and flavonoid contents. **Results:** ME exhibited a strong α -glucosidase inhibition activity (IC_{50} 271.97 μ g/ml) compared to EAE and acarbose (IC_{50} of 1745.05 and 823.99 μ g/ml, respectively). A low dose of ME gave an additive inhibition on α -glucosidase when combined with acarbose. By a kinetic analysis, ME was found to inhibit α -glucosidase in a mixed-type inhibition. Both ME and EAE showed strong antibacterial activities against gram negative and positive bacteria. The strongest inhibition was observed against *C. violaceum* and *S. mutans* for ME (MIC of both 0.387 mg/ml) and *P. aeruginosa* for EAE (MIC 2.938 mg/ml). Both extracts showed weaker antioxidant activities than standards; IC_{50} 664.73 and 4225.40 μ g/ml, respectively. ME was rich in phenolics (277.54 mg GAE/100 g DW), unlike EAE (1.86 mg GAE/100 g DW). **Conclusion:** This study can recommend *B. javanica* seeds as a source for antidiabetic and antibacterial agents. Combination with acarbose may have important role for the treatment of diabetes mellitus.

Key words: Additive inhibition, Antibacterial, *Brucea javanica*, α -glucosidase inhibitor, Mixed type inhibition.

4

Adelina Simamora^{1,*},
Kris Herawan Timotius¹,
Adit Widodo Santoso²

¹Department of Biochemistry, Faculty of Medicine, Krida Wacana Christian University, Jakarta- 11510, INDONESIA.

²Department of Herbal Medicine, Faculty of Medicine, Krida Wacana Christian University, Jakarta- 11510, INDONESIA.

Correspondence

Ms. Adelina Simamora

Department of Biochemistry, Faculty of Medicine, Krida Wacana Christian University, Jakarta, 11510, INDONESIA.

Phone no : +61-21-56942061

E-mail: adelina.simamora@ukrida.ac.id

History

- Submission Date: 05-11-2018;
- Review completed: 03-01-2019;
- Accepted Date: 22-01-2019

DOI : 10.5530/pj.2019.11.76

Article Available online

<http://www.phcogj.com/v11/3>

Copyright

© 2019 Pharmacognosy Journal. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

INTRODUCTION

Diabetes Mellitus (DM) is one of the most common metabolic disorders, which is characterized by high level of postprandial blood glucose (hyperglycemia). This condition may be caused by defect in insulin secretion as a result of impaired pancreatic β -cells, as in type 1 DM. In type 2 DM, this condition may be due to a combination of insulin resistance and insufficient insulin secretion.¹ Around the world, diabetes has affected 346 million people and the number is estimated to increase by twofold in 2030.² Glycemic control is considered an effective therapy in the management of T2DM. Blood glucose is predominantly sourced from enzymatic hydrolysis of dietary polysaccharides by α -amylase and α -glucosidase in the epithelium of small intestine. Accordingly, inhibition of these enzymes can delay the conversion of dietary carbohydrates into glucose, thus lowering postprandial plasma glucose. Therefore, inhibitors that target carbohydrates hydrolyzing enzymes become an important approach in the treatment of hyperglycemia. α -Glucosidase inhibitor drugs such as viglibose, acarbose and miglitol are presently used for treating T2DM. Nevertheless, they are reported to cause unfavorable side effects associated with gastrointestinal

complications such as loose stools, flatulence and abdominal discomfort.³ In view of this, finding new α -glucosidase inhibitors with minimal adverse side effects is of great importance.

Plant materials are good source for drugs, in particular α -glucosidase inhibitors.⁴ *Brucea javanica* (L.) Merr which belongs to *Brucea* genus and Simaroubaceae family, is an evergreen shrub of about 3 m high. It is widely distributed in Asia Pacific regions such as Indonesia, Thailand and Malaysia. The seeds from *B. javanica* have been used traditionally for treating diabetes mellitus.⁵

B. javanica has been previously studied for their biological activities, such as antimalarial,⁶ anti-protozoal,⁷ anti-inflammatory,⁸ antileukemic⁹ and anthelmintic.¹⁰ Some chemical compounds have been reported to be isolated from this plant,¹¹ including quassionoid which is known to have various pharmacological properties such as antitumor and anticancer.¹²⁻¹⁴

To date, only a few studies reported the antidiabetic activities of this plant, such as inhibition activity of the seed extracts against glycogen phosphorylase¹⁵

4

Cite this article: Simamora A, Timotius KH, Santoso AW. Antidiabetic, Antibacterial and Antioxidant Activities of Different Extracts from *Brucea javanica* (L.) Merr Seeds. Pharmacogn J. 2019;11(3):479-85.

and *in vivo* studies using animal models.¹⁶ Nonetheless, there is no report available on *in vitro* inhibitory properties of the seed extract against α -glucosidase. In addition, the mechanism underlying the inhibition and their interaction with the prescribed synthetic inhibitor acarbose is still unknown. Therefore, the study aimed to investigate α -glucosidase inhibition properties of *B. javanica* seeds extracts by *in vitro* enzymatic experiments, including their combination effect with acarbose. The present study extends the investigation by examining the antibacterial and antioxidant potentials of the extracts.

2 MATERIALS AND METHODS

Chemicals and plant material

All solvents and chemicals were analytical grade. 2,2-Diphenyl-1-picryl-hydrazyl (DPPH), 3,5-dinitro-4-butyl-4-hydroxytoluene (BHT), Folin-Ciocalteu's phenol reagent, *p*-nitrophenyl- α -D-glucopyranoside and α -glucosidase (EC 3.2.1.20) from *Saccharomyces cerevisiae* were obtained from Sigma-Aldrich (St. Louis, MO, USA). Rutin and Gallic Acid (GA) were purchased from Santa Cruz Biotechnology (Dallas, USA). Acarbose was purchased from United States Pharmacopeia (Rockville, USA).

The seeds of *B. javanica* were collected from Padang, West Sumatera, Indonesia in March 2016. The plant was identified by one of the authors (KHT). Thoroughly washed seeds were air-dried in the shade and ground to a coarse powder. It was then kept in an air-tight container at 4°C until further used.

Seed extraction

The powder seeds were macerated at ambient temperature overnight successively with n-hexane, ethyl acetate and methanol. Each extract was filtered and dried by vacuum evaporator. The Ethyl Acetate (EAE) and Methanol Extracts (ME) were kept at 4°C in darkness until further use.

Determination of total phenolic content

A colorimetric method using Folin-Ciocalteu reagents was used to estimate the total phenolics of the seed extract, based on previously described procedure.¹⁷ A gallic acid (12.5 – 200 µg/mL) standard curve was obtained with a linear coefficient value of 0.9976 and was used for the estimation of phenolic content of the seed extracts. Data were presented as mg gallic acid equivalent (mg GAE)/100 g dry weight of biomass.¹⁸

Determination of total flavonoid content

The total flavonoid content of the seed extracts was estimated based on aluminum chloride method with slight modification.¹⁹ A rutin (3.2 – 500 µg/mL) standard curve was obtained with a linear coefficient value of 0.9999. This was then used for the estimation of total flavonoid content of the extracts. Data were presented as mg rutin equivalent (mg RE)/100 g of dry weight of biomass.

Determination of α -glucosidase inhibitory activity

Evaluation of inhibitory activity of the seed extracts against α -glucosidase was performed based on our procedure.²⁰ Each extract was diluted to obtain an appropriate dilution series of different concentrations. Into each sample (50 µL) was added DMSO (5 µL), phosphate buffer (45 µL, 50 mM, pH 6.8) and α -glucosidase (50 µL, 0.5 unit/mL). The solution was kept at 37°C for 5 min, after which substrate (100 µL, 1 mM *p*-nitrophenyl- α -D-glucopyranoside) was added to start the reaction. The reaction mixture was then incubated at 37°C for 20 min and stopped by the addition of 750 µL Na₂CO₃ (100 mM). Absorbance was read at 405 nm on a spectrophotometer (Biochrom Libra S-22, Germany). The absorbance of each sample was corrected with blank sample. Acarbose

was used as a positive control of α -glucosidase inhibitor. The percentage (%) of inhibition was calculated as follows:

$$\alpha - \text{Glucosidase inhibition (\%)} = \frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \times 100\%$$

Where: A control is absorbance of control and A sample is absorbance of sample. The IC₅₀ value was described as the extract concentration needed to inhibit 50% of α -glucosidase activity under the assay condition. All measurements were performed in triplicates.

Determination of combined inhibition effect of acarbose and extract on α -glucosidase

Drug combination method developed by Chou-Talalay was used to evaluate combined inhibition effect of acarbose and the most active extract on α -glucosidase.²¹ It required for the inhibitors to be used individually and in combination at different ratios of IC₅₀ obtained from the α -glucosidase assay. α -Glucosidase (0.5 U/mL) and pNPG (1 mM) were used for the assay. The % of inhibition data were then treated using Compusyn® software to calculate Combination Index (CI) for each individual or combined inhibitors. Based on the CI values, drug combinations were considered of having synergistic effects when CI values < 0.9, additive effects CI = 0.9 – 1.1 or antagonistic effects CI > 1.1.

Determination of α -glucosidase inhibition mode

The inhibition mode of the most active extract was determined by performing an enzyme kinetic assay based on previous method.²² In this assay, substrate (*p*-nitrophenyl α -D-glucopyranoside) was prepared at increasing concentrations (0.25 – 1.25 mM). The solution was incubated with α -glucosidase (0.5 U/mL) in the absence and presence of extract (inhibitor) at different concentrations in phosphate buffer (50 mM pH 6.8). Double reciprocal plots of 1/[S] and 1/[V] were generated based on the Lineweaver-Burk method to describe the inhibition mode of the extract.

The Lineweaver-Burk graph was further processed to obtain the values of K_i and K_i'. The K_i value which was the dissociation constant of inhibitor binding to α -glucosidase was obtained from the secondary replot of the slope vs [I] of the Lineweaver-Burk graph. The K_i' value which is the dissociation constant for inhibitor binding to enzyme-substrate (ES) complex was obtained from secondary replot of intercept vs [I].²³

DPPH radical scavenging activity

B. javanica seed extracts was evaluated for their antioxidant activities by measuring the effect of ME and EAE on DPPH radicals, according to a method previously described.²⁴ Into 1 ml of DPPH solution (0.6 mM in ethanol) was added 3 mL of extract or standard solutions (Ascorbic acid and BHT) at different concentrations (10 – 100 µg/mL). The reaction was incubated at ambient temperature in darkness for 30 min. The absorbance was measured at 517 nm with spectrophotometer (Bio Chrom Libra S22). Ethanol (3 mL) in place of extract is used as control. The percentage inhibition activity was determined using the following equation:

$$\text{DPPH inhibition (\%)} = \frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \times 100\%$$

Where: A control is absorbance of control solution and A sample is absorbance of sample. The IC₅₀ value was concentration of extract needed to scavenge 50% of DPPH radicals under the assay condition. All measurements were performed in triplicates.

Antibacterial activity: Agar well diffusion

Evaluation of antibacterial activity of ME and EAE was conducted by an agar well diffusion method on eight bacterial strains i.e. *Staphylococcus aureus* ATCC 25923, *Staphylococcus aureus* COWAN I, *Staphylococcus epidermidis* ATCC 12228, *Chromobacterium violaceum* WT ATCC 12472, *Escherichia coli* ATCC 25922, *Escherichia coli* ATCC 35218, *Pseudomonas aeruginosa* PAO I and *Streptococcus mutans* ATCC 14721. Mueller-Hinton Agar (OXOID CM0337) was used as growth medium. Bacterial suspension (100 µL, McFarland 0.5) was spread on the agar plate. The well diameter was 0.5 cm and filled with 20 µL of the extracts. The incubation was carried out at 37°C for 24 h. The inhibition zone was observed and recorded.

Determination of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)

The MIC and MBC of ME and EAE were determined based on a microwell dilution method in a 96-well plate against the selected bacterial strains. The initial solution was prepared by dissolving each extract in DMSO. This was then followed by preparing serial dilution to give different concentrations. Negative controls were prepared containing bacteria without antimicrobial solution. The bacterial toxicity of the solvent was also evaluated. MIC was determined as a concentration to inhibit the bacterial growth after 24 h. MIC was observed as the last dilution showing no apparent growth (turbidity).

For the determination of MBC, a 10 µL of the culture medium was aspirated from each tube (in the MIC test) that showed no noticeable growth and sub-culturing it on fresh Mueller-Hinton Agar. This was then incubated at 37°C for 24 h. MBC was determined as the lowest concentration of the extracts required to kill the bacterium. MBC was identified as the least concentration showing no noticeable growth on the agar subculture.

Statistical analysis

Experiments in this study were conducted in three replicates. Results were expressed as mean ± Standard Deviation (SD). IC₅₀ values and the kinetics of enzymatic reactions were determined using regression method.

RESULTS**Extract yields, total phenolic and flavonoid contents**

The yields from methanol and ethyl acetate extractions were 27.33% and 1.90% (w/w), respectively. The total phenolic contents in ME and EAE were 277.54 and 1.86 mg GAE/100 g DW (dried weight of biomass), respectively. The total flavonoid contents in ME and EAE were 124.90 and 0.08 mg RE/100 g DW.

The yields are in agreement to those obtained from the previous study by Ablat et al.¹⁵ who reported the lowest yield for ethyl acetate seed extract compared to more polar solvents such as water which obtained the highest yield. However, the same author obtained high contents in phenolics and flavonoids in Ethyl Acetate Extract (EAE) compared to Methanol Extract (ME). This can be attributed to the difference in extraction methods used.

α-Glucosidase inhibition activity

To evaluate the antidiabetic potential of *B. javanica* seeds, the extracts were assessed for their inhibition activity on α-glucosidase. Acarbose, a prescribed synthetic inhibitor, was selected as a positive control to compare the inhibitory activities of the samples. Firstly, ME and EAE were tested for their inhibition activity against α-glucosidase in dose-dependent experiments. Both ME and EAE inhibited α-glucosidase in a concentration dependent manner (Table 1). When the concentration of the extracts increased, the enzyme activity was decreased rapidly. The

12

Table 1: α-Glucosidase inhibitory activities of *B. javanica* seed extracts and acarbose (mean ± SD).

	Concentration (µg/ml)	Inhibition (%)	IC ₅₀ (µg/ml)
ME ^a	98.95	20.08 ± 8.09	271.97 ± 24.99
	197.90	35.49 ± 4.05	
	296.85	63.54 ± 7.18	
	395.80	69.11 ± 9.91	
EAE ^b	376.12	3.34 ± 1.58	1745.05 ± 30.34
	752.24	19.55 ± 8.22	
	1504.48	39.29 ± 8.89	
	1880.60	56.07 ± 3.72	
Acarbose			823.99 ± 0.06

^aME: methanol extract.

^bEAE: ethyl acetate extract.

1

Table 2: CI (Combination index) values of the combined inhibitory activities of acarbose and methanol extracts of *B. javanica* against α-glucosidase.

Acarbose IC ₅₀ value ratio	<i>B. javanica</i> IC ₅₀ value ratio				
	0.25	0.50	0.75	1.00	1.50
0.25	1.03	1.70	1.36	1.49	1.24
0.50	1.14	1.46	1.56	1.68	1.50
0.75	1.13	1.93	1.38	1.53	1.86
1.00	0.98	1.76	1.90	1.39	1.69
1.50	1.06	1.73	1.92	1.56	1.64

Experiments were conducted at different ratio of their IC₅₀ values. Values of 0.25, 0.5, 1 and 1.5 represents 0.25, 0.5, 1 and 1.5-fold of their IC₅₀ values, respectively.

IC₅₀ values for ME and EAE were 271.97±24.99 and 1745.05±30.35 µg/mL, respectively. The IC₅₀ values for acarbose was 823.99±0.06 µg/mL, which is close to previous reports.²⁵

Combined effect

As suggested in our findings above, ME exerted a strong inhibition on α-glucosidase. Hence, it was of interest to investigate whether acarbose and ME may be able to interact synergistically or additively to inhibit α-glucosidase. Experiments were then conducted in solutions containing ME and acarbose in different combinations and the results were presented in Table 2.

When acarbose at different concentrations (0.25 – 1.5 fold of its IC₅₀) was combined with ME at low concentration (0.25 fold of its IC₅₀), 5 CI values in the range of 0.98 – 1.14 were obtained. These values indicate an additive effect. However, when acarbose (0.25 – 1.5 fold of its IC₅₀) was combined with ME at higher concentrations (0.5 – 1.5 fold of its IC₅₀), CI values were calculated in the range of 1.24 – 1.93, suggesting a moderate antagonism.²¹

Type of α-glucosidase inhibition

To further investigate the mechanism underlying the additive inhibition, the inhibition type of ME on α-glucosidase was examined based on the Lineweaver-Burk double reciprocal plots. Figure 1a exhibited that the inhibition of ME on α-glucosidase produced data lines which intersected at one point at the second quadrant of the graph. This result points to a mixed competitive and noncompetitive mode. Our previous work and




Figure 1: (a) Lineweaver-Burk plot for the inhibition of α -glucosidase by *B. javanica* ME. (b) Secondary replot of slope vs [ME] from the Lineweaver-Burk plot for the determination of K_i ($R^2 = 0.9989$) (c) Secondary replot of Y-intercept vs [ME] from the Lineweaver-Burk plot for the determination of K_i ($R^2 = 0.9051$).

other authors found that acarbose inhibited α -glucosidase in a competitive mode.²⁶

Secondary plots were generated for the calculation of K_i and K_i' . The K_i value which was the dissociation constant of ME binding to α -glucosidase was obtained from the secondary replot of slope vs ME of the Lineweaver-Burk graph. A straight line of $R^2 = 0.9989$ (Figure 1b) and K_i value of 650 $\mu\text{g/mL}$ were obtained. The secondary replots of intercept vs ME obtained a straight line with $R^2 = 0.9051$ and $K_i' = 1710 \mu\text{g/mL}$ (Figure 1c), which is the dissociation constant for ME binding to Enzyme-substrate (ES) complex.

Antioxidant activity

DPPH assay assesses the ability of antioxidant compounds to scavenge free radical by donating a hydrogen radical or transferring an electron to the stable DPPH radical, to become its reduced form DPPH-H. Table 3 presents IC_{50} of *B. javanica* seed extracts and standards. Scavenging

Table 3: DPPH radical scavenging activity of *B. javanica* seed extracts and standards (Mean \pm SD).

	Concentration ($\mu\text{g/ml}$)	Inhibition (%)	IC_{50} ($\mu\text{g/ml}$)
ME	123.69	9.04 \pm 0.21	664.73 \pm 5.09
	247.38	17.01 \pm 0.16	
	494.75	36.26 \pm 0.37	
	618.44	46.42 \pm 0.39	
	742.13	56.82 \pm 0.43	
EAE	1062.50	25.09 \pm 1.20	4225.4 \pm 194.62
	2125.00	42.93 \pm 0.11	
	4250.00	54.60 \pm 1.03	
	8500.00	72.35 \pm 1.03	
	Ascorbic acid		53.24 \pm 0.82
	BHT		21.36 \pm 0.80

activity of both ME and EAE on DPPH radical were evident, with ME showed a stronger activity of more than 6 times stronger than EAE. However, compared to the standards BHT and ascorbic acid, the activities were much weaker.

Antibacterial activity

Antibacterial activity of seed extracts from *B. javanica* were examined against gram positive bacteria (*S. aureus*, *S. epidermidis* and *S. mutans*) and gram-negative bacteria (*E. coli*, *P. aeruginosa* and *C. violacium*) by determining the DIZ, MIC and MBC. The results are presented on Table 4. ME and EAE exhibited notable antibacterial activity against all tested bacteria. In general, ME exhibited stronger activity than EAE with MICs in the range of 0.387 to 3.092 mg/mL and MBC in the range of 0.774 to 6.184 mg/mL. Among different bacteria, gram negative strain *E. coli* and *S. aureus* were found to be the most sensitive to ME (DIZ of 20.75 and 20.20 mm, respectively), whereas *S. aureus* was the most sensitive to EAE (DIZ of 18.90 mm).

DISCUSSION

Natural products from plants are still valuable sources to develop alternative compounds with low side effect but having desirable α -glucosidase inhibition property. Ethnobotanical information indicates that around 400 plant medicines have been used to control hyperglycemia.²⁷ In another report, more than 400 compounds exhibiting inhibitory activity on α -glucosidase have been identified from plant medicine,²⁸ suggesting the potential of medicinal plants to be employed in management of DM. In this direction, we set to investigate the α -glucosidase inhibition properties of *B. javanica*.

In this work, it was found that the seeds of *B. javanica* extracted in methanol was a more effective α -glucosidase inhibitor, three times stronger than the standard acarbose. On the other hand, EAE showed a weaker inhibition activity when compared with acarbose. Association between polyphenolics and inhibition of key enzymes associated with carbohydrate hydrolyzes has been reported previously.²⁹ Further, it has been reported that these secondary metabolites modulate enzymatic breakdown of dietary carbohydrates through binding with α -glucosidase.³⁰⁻³¹ In view of this, the difference in inhibition activities between ME and EAE may be associated with the level of phenolic and flavonoid in the extracts. To the best of our knowledge, the present study is the first time α -glucosidase inhibition activity was reported for *B. javanica*, which may be a new source of inhibitor for T2DM treatment.

Table 4: Antibacterial activity (MIC and MBC) of *B. javanica* seed extracts (mean \pm SD).

	ME			EAE		
	DZI	MIC	MBC	DZI	MIC	MBC
<i>S. aureus</i> 25923	20.2 \pm 0.2	0.003092	0.006184	18.9 \pm 0.3	0.023508	0.047016
<i>S. aureus</i> COWAN 1	10.85 \pm 0.03	0.001546	0.003092	9.65 \pm 0.4	0.011754	0.023508
<i>S. epidermidis</i> 12228	14.65 \pm 0.02	0.001546	0.003092	11.75 \pm 0.5	0.023508	0.047016
<i>Chr. violaceum</i> 12472	8.9 \pm 0.01	0.000387	0.000774	7.8 \pm 0.09	0.023508	0.047016
<i>E. coli</i> 25922	20.75 \pm 0.1	0.001546	0.003092	11.9 \pm 0.1	0.005877	0.011754
<i>E. coli</i> 35218	15.7 \pm 0.3	0.003092	0.006184	9.25 \pm 0.17	0.011754	0.023508
<i>P. aeruginosa</i> PAO 1	15.25 \pm 0.09	0.001546	0.003092	8.02 \pm 0.43	0.002938	0.005876
<i>Str. mutans</i> 14721	8.9 \pm 0.01	0.000387	0.000774	7.95 \pm 0.27	0.005877	0.011754

³⁶ Z: Diameter of inhibition zone.

^a MIC: Minimum inhibitory concentration.

^c MBC: Minimum bactericidal concentration.

In this study it was found that the combination of acarbose at various concentrations and ME at low concentrations exerted an additive effect. This effect indicates that ME may substitute for or be used in conjunction with acarbose in maintaining glycemic control in the management of T2DM. Previous studies have reported combined inhibition of plant extracts and acarbose. For instance, Adisakwattana *et al.*³² reported additive inhibition for combination of cinnamon bark aqueous extracts and acarbose on intestinal α -glucosidase. However, no information available on the range of acarbose dose and the amount of cinnamon extracts concentrations required to produce the effect. In the present study ME and acarbose exhibited additive inhibition at low concentrations and turned into moderate antagonistic effect at higher concentrations.

Long term use of acarbose leads to accumulation of undigested carbohydrates in the colon, which serves as substrates for bacteria fermentation. In consequence of this is troublesome gastrointestinal symptoms such as diarrhea and flatulence.³³ In some cases, long term use of acarbose was reported to cause acute severe hepatotoxicity.³⁴ These side effects tend to increase in higher dose, as in the case of chronic T2DM patients where progressive increase dosages are prescribed. Therefore, reduction of the required dose of acarbose may potentially minimize the multiple adverse effects of this medicine. Furthermore, this may ultimately reduce the cost associated with pharmaceutical DM management. Our study suggests that combinations of acarbose and ME at low dose may serve as a possible combination therapy in preference of only acarbose treatment.

The study found that *B. javanica* seed extract gave a mixed type inhibition of non-competitive and competitive, whereas acarbose inhibited α -glucosidase in a competitive manner. When compared, the K_i value of ME was almost three times more than K_i value, suggesting that binding of ME to the ES was stronger than ME to enzyme alone. This finding indicates that the inhibition of ME was noncompetitive, predominant over competitive. The additive effect exerted from the combination of ME and acarbose is possibly come from the different inhibition binding sites of the two inhibitors at α -glucosidase.

This study also found that the seed extracts showed antioxidant activities. Many studies demonstrated that chronic hyperglycemia as in T2DM induced generation of free radicals through glucolipotoxicity and mitochondrial pathways.³⁵ This condition may cause oxidative damages in cell membrane, leading to the development of various complications in diabetes, including retinopathy, cardiomyopathy, neuropathy and nephropathy.³⁶ In particular, destruction in pancreatic β cells which are known to be vulnerable to free radicals ultimately results in impaired insulin secretion.³⁷ Therefore, antioxidant intake such as from diets may

help protect cells from damages, thus may prevent diabetes and diabetic complications.

The present study also found that ME and EAE exhibited good antibacterial activity. Good antibacterial activity observed for EAE indicates that phenolics and flavonoids may not play major role in inhibiting the tested bacteria. Previously, many bioactive compounds have been identified from different parts of *B. javanica*, such as derivatives of quassinoids that had anticancer and antiprotozoal activities.¹⁴ However, to the extent of our knowledge, only a few studies reported antibacterial activities of *B. javanica*. Previous reports included anti-tuberculosis activity of quassinoids isolated from *B. javanica* seeds³⁸ and an antibacterial peptide, named Brucin isolated from the fruit of *B. javanica* which had specificity for *Streptococcus pyogenes*.³⁹ Thus, the current study evidences the potential of the seeds from *B. javanica* to be explored as antibacterial agent.

CONCLUSION

The present study concludes that methanol seed extract of *B. javanica* has a therapeutic potential better than ethyl acetate extract, thus can be a new potential source for antidiabetic and antibacterial agent. Combination of *B. javanica* seed extract with acarbose is likely to be employed in diabetes therapy, with a possibility of reducing acarbose dosage.

ACKNOWLEDGEMENT

This work was fully supported by the Faculty of Medicine, Krida Wacana Christian University.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ABBREVIATIONS


ME: Methanol Extract; EAE: Ethyl Acetate Extract; GAE: Gallic Acid Equivalence; DPPH: 2,2-diphenyl-1-picryl-hydrazyl.

REFERENCES

- American Diabetes Association. Diagnosis and classification of diabetes mellitus. Diabetes care. 2014;37(Supplement 1):S81-90.
- Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: global estimates of the prevalence of diabetes for 2011 and 2030. Diabetes Res Clin Pract. 2011;94(3):311-21.
- Nyenwe EA, Jenkins TW, Umpierrez GE, Kitabchi AE. Management of type 2 diabetes: evolving strategies for the treatment of patients with type 2 diabetes. Metabolism. 2011;60(1):1-23.
- Choudhury H, Pandey M, Hua CK, Mun CS, Jing JK, Kong L, *et al.* An update on

- natural compounds in the remedy of diabetes mellitus: a systematic review. *J Tradit Complement Med*. 2018;8(3):361-76.
5. Shamsul K, Tajuddin A, Mazina M. *Tumbuhan ubatan tradisional Malaysia*. Institut Biosains, Universiti Pertanian Malaysia Publisher, Malaysia. 2003;170.
 6. Kefe A, Giday M, Mamo H, Erko B. Antimalarial properties of crude extracts of seeds of *Brucea antidysenterica* and leaves of *Ocimum lamiifolium*. *BMC Complement Altern Med*. 2016;16(1):118-26.
 7. Kanyanga CR, Ehata TM, Lumpu NS, Munduku KC, Phuati MA, Kabangu KO, et al. Antiprotozoal and cytotoxic activities and the acute toxicity of extracts from *Brucea sumatrana* Roxb. (*Simaroubaceae*) leaves collected in Mai-Ndombe, Democratic Republic of Congo. *J Pharmacognosy Phytother*. 2015;7(4):35-44.
 8. Huang YF, Zhou JT, Qu C, Dou YX, Huang QH, Lin ZX, et al. Anti-inflammatory effects of *Brucea javanica* oil emulsion by suppressing NF- κ B activation on dextran sulfate sodium-induced ulcerative colitis in mice. *J Ethnopharmacol*. 2017;198:389-98.
 9. Lee K-H, Hayashi N, Okano M, Nozaki H, Ju-Ichi M. Antitumor agents, 65. Brusatol and cleomiscosin-A, antileukemic principles from *Brucea javanica*. *J Nat Prod*. 1984;47(3):550-1.
 10. Wang Y, Wu ZF, Wang GX, Wang F, Liu YT, Li FY, et al. *In vivo* anthelmintic activity of bruceine A and bruceine D from *Brucea javanica* against *Dactylogyrus intermedius* (*Monogenea*) in goldfish (*Carassius auratus*). *Vet Parasitol*. 2011;177(1-2):127-33.
 11. Dong SH, Liu J, Ge YZ, Dong L, Xu CH, Ding J, et al. Chemical constituents from *Brucea javanica*. *Phytochemistry*. 2013;85:175-84.
 12. Yan Z, Guo G-f, Zhang B. Research of *Brucea javanica* against cancer. *Chin J Integr Med*. 2017;23(2):153-60.
 13. Zhao L, Li C, Zhang Y, Wen Q, Ren D. Phytochemical and Biological Activities of an Anticancer Plant Medicine: *Brucea javanica*. *Anti-Cancer Agents in Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Cancer Agents)*. 2014;14(3):440-58.
 14. Zhao L, Li C, Zhang Y, Wen Q, Ren D. Phytochemical and biological activities of an anticancer plant medicine: *Brucea javanica*. *Anticancer Agents Med Chem*. 2014;14(3):440-58.
 15. Ablat A, Mohamad J, Awang K, Shilpi JA, Arya A. Evaluation of antidiabetic and antioxidant properties of *Brucea javanica* seed. *The Scientific World Journal*. 2014;2014:1-8.
 16. Noorshahida A, Wong TW, Choo CY. Hypoglycemic effect of quassinoids from *Brucea javanica* (L.) Merr (*Simaroubaceae*) seeds. *J Ethnopharmacol*. 2009;124(3):586-91.
 17. Singleton V, Rossi JA. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am J Enol Vitic*. 1965;16(3):144-58.
 18. Lee S, Mediani A, Nur AA, Azliana A, Abas F. Antioxidant and α -glucosidase inhibitory activities of the leaf and stem of selected traditional medicinal plants. *Int Food Res J*. 2014;21(1):379-86.
 19. Kamtekar S, Keer V, Patil V. Estimation of phenolic content, flavonoid content, antioxidant and α amylase inhibitory activity of marketed polyherbal formulation. *Journal of Applied Pharmaceutical Science*. 2014;4(9):61.
 20. Simamora A, Paramita L, Mohamad HNAB, Santoso AW, Timotius KH. *In vitro* antidiabetic and antioxidant activities of aqueous extract from the leave and fruit of *Psidium guajava* L. *Indones Biomed J*. 2018;10(2):156-64.
 21. Chou TC. Theoretical basis, experimental design and computerized simulation of synergism and antagonism in drug combination studies. *Pharmacol Rev*. 2006;58(3):621-81.
 22. Timotius KH, Simamora A, Santoso AW. Chemical characteristics and *in vitro* antidiabetic and antioxidant activities of *Premna serratifolia* L. leaf infusion and decoction. *Pharmacog J*. 2018;10(6):1114-8.
 23. Adisakwattana S, Charoenlerkul P, Yibchok-anun S. α -Glucosidase inhibitory activity of cyanidin-3-galactoside and synergistic effect with acarbose. *J Enzyme Inhib Med Ch*. 2009;24(1):65-9.
 24. Simamora A, Adit WS, Kris HT. Bioactivities of Methanol and Ethyl Acetate Mace Extracts of *Myristica fragrans* Houtt. *Pharmacognosy Communications*. 2018;8(3).
 25. Ali RB, Atangwho IJ, Kuar N, Ahmad M, Mahmud R, Asmawi MZ. *In vitro* and *in vivo* effects of standardized extract and fractions of *Phaleria macrocarpa* fruits pericarp on lead carbohydrate digesting enzymes. *BMC Complement Altern Med*. 2013;13(1):39.
 26. Zhang BW, Sang YB, Sun WL, Yu HS, Ma BP, Xiu ZL, et al. Combination of flavonoids from *Oroxylum indicum* seed extracts and acarbose improves the inhibition of postprandial blood glucose: *In vivo* and *in vitro* study. *Biomed Pharmacother*. 2017;91:890-8.
 27. Bailey CJ, Day C. Traditional plant medicines as treatments for diabetes. *Diabetes Care*. 1989;12(8):553-64.
 28. Yin Z, Zhang W, Feng F, Zhang Y, Kang W. α -Glucosidase inhibitors isolated from medicinal plants. *FSHW*. 2014;3(3-4):136-74.
 29. Shobana S, Sreerama Y, Malleshi N. Composition and enzyme inhibitory properties of finger millet (*Eleusine coracana* L.) seed coat phenolics: mode of inhibition of α -glucosidase and pancreatic amylase. *Food Chem*. 2009;115(4):1268-73.
 30. Pantidos N, Boath A, Lund V, Conner S, McDougall GJ. Phenolic-rich extracts from the edible seaweed, *ascophyllum nodosum*, inhibit α -amylase and α -glucosidase: potential anti-hyperglycemic effects. *J Func Foods*. 2014;10:201-9.
 31. Tadera K, Minami Y, Takamatsu K, Matsuoka T. Inhibition of α -glucosidase and α -amylase by flavonoids. *J Nutr Sci Vitaminol*. 2006;52(2):149-53.
 32. Adisakwattana S, Lerdswankij O, Poputtachai U, Minipun A, Suparprom C. Inhibitory activity of cinnamon bark species and their combination effect with acarbose against intestinal α -glucosidase and pancreatic α -amylase. *Plant Foods Hum Nutr*. 2011;66(2):143-8.
 33. Kim JG, Jo SH, Ha KS, Kim SC, Kim YC, Apostolidis E, et al. Effect of long-term supplementation of low molecular weight chitosan oligosaccharide (GO2KA1) on fasting blood glucose and HbA1c in db/db mice model and elucidation of mechanism of action. *BMC Complement Altern Med*. 2014;14(1):272.
 34. Carrascosa M, Pascual F, Aresti S. Acarbose induced acute severe hepatotoxicity. *The Lancet*. 1997;349(9053):698-9.
 35. Mohora M, Greabu M, Musculel C, Duta C, Totan A. The Sources and the targets of oxidative stress in the etiology of diabetic complications. *Rom J Biophys*. 2007;17(2):63-84.
 36. Yan LJ. Pathogenesis of chronic hyperglycemia: from reductive stress to oxidative stress. *J Diabetes Res*. 2014;2014.
 37. Li N, Frigerio F, Maechler P. The sensitivity of pancreatic β -cells to mitochondrial injuries triggered by lipotoxicity and oxidative stress. *Portland Press Limited*. 2008.
 38. Rahman S, Fukamiya N, Okano M, Tagahara K, Lee KH. Anti-tuberculosis activity of quassinoids. *Chem Pharm Bull*. 1997;45(9):1527-9.
 39. Sornwatana T, Roytrakul S, Wetprasit N, Ratanapo S. Brucin, an antibacterial peptide derived from fruit protein of Fructus *Bruceae*, *Brucea javanica* (L.) Merr. *Lett Appl Microbiol*. 2013;57(2):129-36.

GRAPHICAL ABSTRACT



SUMMARY

- The seeds of *Brucea javanica* was known as herbal material for anticancer, antimalarial and anti-amoebs. Its antidiabetic properties are not yet well understood. Our study was the first to report its inhibition activity against α -glucosidase, a key enzyme in carbohydrate metabolism. Methanol extract of the seeds is found to be a potent inhibitor. Its combination with acarbose gives rise to an additive effect. In addition, both methanol and ethyl acetate extracts were found to have notable anti-bacterial activities. This study recommends *B. javanica* as a new source for antidiabetic agent

ABOUT AUTHORS



Adelina Simamora, MSc (Pharm) is a researcher in the centre for herbal medicine studies and a lecturer in Biochemistry at the Faculty of Medicine Krida Wacana Christian University. At present, her research focus on the bioactive potential of medicinal plants, with special interest in various enzyme inhibition properties. Currently, her work involves various Indonesian medicinal plants and inhibition properties on α -glucosidase, xanthine oxidase, and lipase. Her interest also spans to transition and rare earth metals and their application as sensor in biological fluid.



Dr Kris Herawan Timotius is a lecturer in microbiology and biochemistry at the Faculty of Medicine Krida Wacana Christian University. His research focus on herbal medicine including natural antibiotics, anti-quorum sensing, antibiofilm, enzyme therapy, and anti enzyme activities such as alpha glucosidase and xanthine oxidase.



Adit Widodo Santoso, BSc: A researcher in the centre for herbal medicine studies. His interest is mainly in the areas of microbiological activities of variety of plant species, including antibiofilm and anti-quorum sensing activities. Currently, he is focusing on bioautography involving thin layer chromatography and western blot. He is also interested in zymography studies and their applications.

Cite this article: Simamora A, Timotius KH, Santoso AW. Antidiabetic, Antibacterial and Antioxidant Activities of Different Extracts from *Brucea javanica* (L.) Merr Seeds. Pharmacog J. 2019;11(3):479-85.

ORIGINALITY REPORT

23%

SIMILARITY INDEX

17%

INTERNET SOURCES

19%

PUBLICATIONS

12%

STUDENT PAPERS

PRIMARY SOURCES

1

www.foodandnutritionjournal.org

Internet Source

5%

2

pdfs.semanticscholar.org

Internet Source

3%

3

Sirichai Adisakwattana, Piyawan Charoenlertkul, Sirintorn Yibchok-anun. " α -Glucosidase inhibitory activity of cyanidin-3-galactoside and synergistic effect with acarbose", Journal of Enzyme Inhibition and Medicinal Chemistry, 2008

Publication

2%

4

www.phcogj.com

Internet Source

1%

5

Adelina Simamora, Lusia Paramita, Nur Azreen, Adit Widodo Santoso, Kris Herawan Timotius. "In Vitro Antidiabetic and Antioxidant Activities of Aqueous Extract from the Leaf and Fruit of *Psidium guajava* L.", The Indonesian Biomedical Journal, 2018

Publication

1%

6	www.tandfonline.com Internet Source	1 %
7	www.hindawi.com Internet Source	1 %
8	Ganiyu Oboh, Opeyemi Babatunde Ogunsuyi, Mariam Damilola Ogunbadejo, Stephen Adeniyi Adefegha. "Influence of gallic acid on α -amylase and α -glucosidase inhibitory properties of acarbose", Journal of Food and Drug Analysis, 2016 Publication	1 %
9	Submitted to Academic Library Consortium Student Paper	1 %
10	www.mdpi.com Internet Source	<1 %
11	Submitted to Universitas Indonesia Student Paper	<1 %
12	ADELINA SIMAMORA, ADIT WIDODO SANTOSO, KRIS HERAWAN TIMOTIUS. " α -Glucosidase Inhibitory Effect of Fermented Fruit Juice of Morinda Citrifolia L and Combination Effect with Acarbose", Current Research in Nutrition and Food Science Journal, 2019 Publication	<1 %
13	www.thieme-connect.com Internet Source	<1 %

14	Bo-wei Zhang, Yuan-bin Sang, Wen-long Sun, He-shui Yu, Bai-ping Ma, Zhi-long Xiu, Yue-sheng Dong. "Combination of flavonoids from Oroxylum indicum seed extracts and acarbose improves the inhibition of postprandial blood glucose: In vivo and in vitro study", Biomedicine & Pharmacotherapy, 2017 Publication	<1 %
15	Submitted to University of Rajshahi Student Paper	<1 %
16	studentsrepo.um.edu.my Internet Source	<1 %
17	Nusaibah Zahratunnisa, Berna Elya, Arikadia Noviani. "Inhibition of Alpha-Glucosidase and Antioxidant Test of Stem Bark Extracts of Garcinia fruticosa Lauterb", Pharmacognosy Journal, 2017 Publication	<1 %
18	www.frontiersin.org Internet Source	<1 %
19	www.scialert.net Internet Source	<1 %
20	www.scipharm.at Internet Source	<1 %
21	Manzoor, Maleeha, Farooq Anwar, Nazamid	

Saari, and Muhammad Ashraf. "Variations of Antioxidant Characteristics and Mineral Contents in Pulp and Peel of Different Apple (*Malus domestica* Borkh.) Cultivars from Pakistan", *Molecules*, 2012.

Publication

<1 %

22

Li, Y.. "Punica granatum flower extract, a potent α -glucosidase inhibitor, improves postprandial hyperglycemia in Zucker diabetic fatty rats", *Journal of Ethnopharmacology*, 20050603

Publication

<1 %

23

Choi, Chun Whan, Yeon Hee Choi, Mi-Ran Cha, Dae Seok Yoo, Young Sup Kim, Gyu Hwan Yon, Kyung Sik Hong, Young Ho Kim, and Shi Yong Ryu. "Yeast α -Glucosidase Inhibition by Isoflavones from Plants of Leguminosae as an in Vitro Alternative to Acarbose", *Journal of Agricultural and Food Chemistry*, 2010.

Publication

<1 %

24

www.nature.com

Internet Source

<1 %

25

İbrahim Kivrak. "Analytical Methods Applied to Assess Chemical Composition, Nutritional Value and In Vitro Bioactivities of *Terfezia olbiensis* and *Terfezia claveryi* from Turkey", *Food Analytical Methods*, 2014

Publication

<1 %

26	Ganiyu Oboh. "Effect of Combination on the Antioxidant and Inhibitory Properties of Tropical Pepper Varieties Against α -Amylase and α -Glucosidase Activities <i>In Vitro</i> ", Journal of Medicinal Food, 06/11/2011 Publication	<1 %
27	Submitted to Institute of Chemical Technnology, Mumbai Student Paper	<1 %
28	Submitted to Liverpool John Moores University Student Paper	<1 %
29	Submitted to Chulalongkorn University Student Paper	<1 %
30	journals.plos.org Internet Source	<1 %
31	Submitted to Universiti Putra Malaysia Student Paper	<1 %
32	Submitted to Middle East Technical University Student Paper	<1 %
33	www.wjgnet.com Internet Source	<1 %
34	Submitted to University of Abertay Dundee Student Paper	<1 %
35	Chandrasekara, Anoma, and Fereidoon Shahidi.	<1 %

"Inhibitory Activities of Soluble and Bound Millet Seed Phenolics on Free Radicals and Reactive Oxygen Species", Journal of Agricultural and Food Chemistry, 2011.

Publication

36

Submitted to Cyprus International University

Student Paper

<1%

37

www.sigadmin.ufpi.br

Internet Source

<1%

Exclude quotes On

Exclude bibliography On

Exclude matches

< 10 words