

A meta-analysis

by Dame Elysabeth

Submission date: 05-Sep-2023 10:51AM (UTC+0700)

Submission ID: 2157981053

File name: A_meta-analysis.pdf (2.64M)

Word count: 8114

Character count: 45963

ORIGINAL MANUSCRIPT

A meta-analysis of studies of the effects of case management intervention for stroke survivors across three countries

Ita Daryanti Saragih MSN, RN¹ | Ice Septriani Saragih MSN, RN² |
Dame Elysabeth Tuty Arna Uly Tarihoran MSN, RN^{3,4} | Sapna Sharma MSN, RN¹ |
Fan-Hao Chou PhD, RN, FAAN¹

¹College of Nursing, Kaohsiung Medical University, Kaohsiung, Taiwan

²STikes Santa Elisabeth Medan, Medan, Indonesia

³School of Nursing, Universitas Kristen Wacana Jakarta, Indonesia

⁴School of Nursing Faculty of Medical and Health Sciences, University of Auckland, Auckland, New Zealand

Correspondence

Fan-Hao Chou, PhD, RN, FAAN, College of Nursing, Kaohsiung Medical University, 100, Shih-Chuan 1st Road, Kaohsiung 80708, Taiwan.

Email: fanhao@kmu.edu.tw

Abstract

Purpose: A stroke survivor who is discharged to home care faces many challenges during the transition from the hospital to home, including managing their care at home and engaging in rehabilitation and recovery. Case management was developed to ease the transition from hospital to home and help people with stroke cope with their care management challenges. However, the effects of case management intervention remain inconclusive. Case management was designed to direct care and may represent a novel method for reducing the burden of care. This study was designed to evaluate the effects of case management interventions on mental health outcomes, activities of daily living capacity, physical function, and social function among stroke survivors.

Design: Systematic review and meta-analysis.

Methods: This study included studies examining the impacts of case management interventions for stroke survivors that were published in English and identified by searching eight databases, from database inception to February 20, 2022. Standardized mean differences (SMDs) with 95% confidence intervals (CIs) were used to pool effect sizes using a random-effects model (in Stata 16.0). The revised Cochrane risk-of-bias tool for randomized trials (RoB-2) was used to assess the methodological quality of each study.

Findings: The inclusion criteria were satisfied by eight studies (including a total of 1119 stroke survivors). Case management had positive effects on mental health (SMD: 0.26; 95% CI: 0.07 to 0.45, $p = 0.001$) and activities of daily living (SMD: 0.68; 95% CI: 0.037 to 0.99, $p < 0.001$). However, no significant effects were observed for either physical function or social function.

Conclusion: Case management appears to enhance the mental health and activities of daily living among stroke survivors.

Clinical Relevance: Case management interventions hold promise as efficient, cost-effective, and accessible strategies to positively influence care for stroke survivors. This intervention strategy could be applied to the hospital-to-home transition to guide care among this population.

KEYWORDS

case management intervention, mental health, meta-analysis, rehabilitation, stroke survivors

INTRODUCTION

Stroke is the leading cause of death and disability worldwide (Campbell & Khatri, 2020) and is associated with considerable personal, family, and societal burdens, including high economic and social costs (Chen, Wei, et al., 2021; Wang et al., 2021). In 2019, approximately 101.5 million people were estimated to be living with stroke, and stroke was responsible for over 6.6 million deaths worldwide (Alonzo et al., 2021; Virani et al., 2021). Currently, more than 80 million stroke survivors are considered disabled, and this population is estimated to increase over the coming decade, with an expected 50% increase in new stroke survivors each year (Duncan et al., 2021; Mendis et al., 2015).

Stroke episodes are characterized by symptoms including sudden numbness or weakness, confusion, difficulty speaking, difficulty seeing with one or both eyes, and severe headaches without an underlying cause (Centers for Disease Control and Prevention, 2020; Rodgers & Price, 2017). The sudden changes in the condition experienced by post-stroke survivors lead to a variety of obstacles and challenges when performing activities of daily living and can include emotional and behavioral changes (Khazaal et al., 2021; Towfighi et al., 2017). As a consequence of the changes associated with stroke, survivors often experience declines in social engagement, work participation, family engagement, community engagement, and recreational activities (Espenberger et al., 2021; Tse et al., 2017). Approximately 1 in 3 post-stroke survivors (Guo et al., 2021) are estimated to develop depression symptoms in response to their inability to cope with daily stressors and the loss of ability to function independently (Ariful Islam et al., 2016; Saragih, Tarigan, Batubara, Tzeng, & Lin, 2021a; Towfighi et al., 2017).

Most stroke survivors are eventually discharged from the hospital, and the hospital-to-home transition introduces several hurdles to recovery and rehabilitation (Clery et al., 2020; Lindblom et al., 2020; Reeves et al., 2019). Caregivers for stroke survivors face difficult challenges during the transition from hospital to home, including emotional, social, and physiological concerns (Grant et al., 2014; Harford et al., 2019). The transition from the hospital to the home represents a critical window during which stroke survivors and their caregivers must foster self-management skills and establish the groundwork necessary to achieve long-term recovery and rehabilitation following a stroke (Broderick & Abir, 2015; Lin, Xiao, & Chamberlain, 2020a; Reeves et al., 2017b). Prominent challenges faced during the hospital-to-home transition include identifying methods to assist stroke survivors in the management of various factors that affect their mental health, emotional status, and ability to perform activities of daily living (Camicia et al., 2016; Chen et al., 2020; Clarke & Forster, 2015).

One strategy used to improve care for this population is case management, with the goal of integrating services by providing care management to help patients with long-term conditions meet and cope with the challenges associated with recovery and rehabilitation (Mountain et al., 2020; Reeves et al., 2017a). Case management is defined as "a health-care process in which a professional assists a

patient with planning, facilitation, care coordination, and advocacy to achieve the best possible healthcare and psychosocial outcomes while also encouraging cost-effectiveness" (Hudon et al., 2019). Case management can assist in directing therapy depending on the circumstances and requirements of those receiving it (Saragih, Tonapa, Lin, & Lee, 2021b). Case management represents a team endeavor in which case managers provide health and human services such as screening, assessment, risk stratification, planning, administration, coordination, monitoring, transitioning and communicating post-transition, and evaluating the care required to meet patient's needs through extensive and integrated health care procedures (Fabbri et al., 2017; Lukersmith et al., 2016; Woodward & Rice, 2015). Consequently, case management interventions are generally considered compelling techniques for decreasing medical care expenses and increasing health care utilization (Brazle, 2019; Joo & Liu, 2017).

The impacts of case management interventions on stroke survivors have been examined in several studies. A randomized controlled trial demonstrated that a 3-month case management intervention decreased depression in a population in the United States ($p < 0.001$) and increased self-care adherence ($p < 0.05$) (Claiborne, 2006); an 8-week case management intervention, however, was found to have no effect on depression ($p = 0.97$) (Reeves et al., 2019). Another 6-month case management program in Canada resulted in a significant improvement in physical function ($p < 0.05$) (Mayo et al., 2009); however, another study conducted over a 12-month period showed no improvement in physical ($p = 0.24$) or social function ($p = 0.24$) (Markle-Reid et al., 2011). Furthermore, a substantial increase in the ability to perform activities of daily living was observed following an 8-week case management intervention performed in the United States among stroke survivors ($p = 0.007$) (Mayo & Scott, 2011). However, no changes in activities of daily living were observed in the United States following a 6-month intervention ($p = 0.47$) (Mayo et al., 2009).

Although these studies have increased our understanding of the potential advantages associated with the use of case management interventions among stroke survivors, the outcomes of this intervention remain uncertain. However, identifying an appropriate intervention that is able to address the physical and social functional limitations, in addition to the mental health and emotional problems experienced by stroke survivors, remains necessary to improve their quality of life. To provide a more comprehensive view of this phenomenon, data regarding the consequences of case management interventions among stroke survivors should be collected and analyzed. The goal of this systematic review and meta-analysis was to assess the effects of case management interventions on stroke survivors' mental health, ability to perform activities of daily living, and physical and social functions.

DESIGN

This review was registered with the International Prospective Register of Systematic Reviews (PROSPERO: CRD42021283266).

This investigation was performed in accordance with the most recent Preferred Items for Systematic Reviews and Meta-Analyses (PRISMA) criteria (Page et al., 2021) (Appendix S1).

81 MATERIAL AND METHODS

Search strategy

One reviewer conducted a systematic literature search to identify literature indexed from database inception to February 20, 2021, using eight databases: Academic Search Complete, Cochrane library, CINAHL, EMBASE, MEDLINE, PubMed, OVID (UpToDate), and Web of Science. Medical Subject Heading terms were utilized in the search, including "stroke patients" OR "post-stroke" OR "stroke survivors" OR "after stroke" AND "case management" OR "case management model" OR "case management nursing" OR "case managed" AND "randomized controlled trial" OR "randomized controlled study" OR "randomized" OR "clinical trials" OR "allocation random." Appendix S1 contains an overview of the search strategy.

31 Eligibility criteria

The PICOS (Population, Intervention/issue of interest, Comparison, Outcome, and Study design) approach was employed to examine the inclusion criteria (Liberati et al., 2009). The inclusion criteria were studies with stroke survivors, intervention groups that included case management interventions, randomized controlled trials, and original research. Studies that did not match the PICOS requirements, studies that did not have full text available, and studies that did not adequately describe their research techniques were excluded.

Study selection and data extraction

Two reviewers assessed 50 relevance of identified studies based on the title and abstract and removed those that did not meet the PICOS criteria (Amir-Behghadami & Janati, 2020). A full-text screening of 11 remaining studies was performed, and those that satisfied the inclusion criteria were included in the study. Following the selection of the final papers, two independent reviewers extracted crucial data items, including the author, the year of publication, the country the study was performed in, the research design, the interventionist, the intervention setting, the demographic data of the participants (e.g., total participants, number of women, and age), intervention details (e.g., intervention types compared between groups, frequency and period of intervention, and follow-up), and outcomes. Any disagreements that developed during the process were resolved through a consensus meeting with a third reviewer.

28 Quality assessment

The methodological quality of the selected studies was assessed by two reviewers using the Cochrane risk-of-bias assessment for randomized trials (RoB-2), and all disagreements were resolved through a consensus meeting with a third reviewer.

Data synthesis and analysis


Because different instruments were used to evaluate the same outcomes across the included studies, the standardized mean difference (SMD) was estimated using a 95% confidence interval (CI) (Liu et al., 2017; Sedgwick & Marston, 2013). The impacts of the intervention on various outcomes were evaluated by integrating data collected using multiple instruments for each study group and comparing the mean and standard deviation between pre- and post-intervention assessments. For both the intervention and control groups, the mean difference and standard deviation were computed, and the SMD effect size (Cohen's d) was calculated (Cahan & Gamliel, 2011; Wilson, 2016).

The existence of heterogeneity within the measured variables, including mental health outcomes, activities of daily living capacity, physical function, and social function, examined among stroke survivors was examined using meta-analysis. For each variable, heterogeneity was assessed using the random-effects model by Q , τ^2 , and I^2 ; I^2 values of 25%, 50%, and 75% indicated low, moderate, and high heterogeneity, respectively (Higgins et al., 2003). The Egger's test and a funnel plot were used to visualize publication bias (Egger et al., 1997). STATA 16.0 was utilized to determine the effect sizes of the variables in the meta-analysis.

18 RESULTS

Study selection

A total of 2093 studies were identified among the eight databases, 666 of which were discarded as duplicates. The remaining 1427 papers were examined based on title and abstract, which resulted in 1403 studies being discarded because they did not match the PICOS criteria, as follows: the population of interest was not the research population ($n = 638$); the case management intervention was not used in the trials ($n = 703$); the study was a review study ($n = 57$); the study was not available in English ($n = 5$). The full texts of the remaining 24 studies were assessed against the eligibility criteria, and 16 were removed because they did not employ case management interventions ($n = 7$), were review studies ($n = 7$), were not published in a peer-reviewed journal ($n = 7$), or were not published in English ($n = 2$). The final qualitative synthesis and meta-analysis included eight studies (Claiborne, 2006; Kummarg et al., 2018; Markle-Reid et al., 2011; Mayo et al., 2008; Mayo et al., 2009; Mayo & Scott, 2011; Reeves et al., 2019; Saal



65 **FIGURE 1** PRISMA flowchart diagram

et al., 2015). ²² Figure 1 depicts the study selection process using a PRISMA flowchart.

Study characteristics

Between 2006 and 2019, eight RCTs ³ were published, including four studies performed in Canada (Markle-Reid et al., 2011; Mayo et al., 2008; Mayo et al., 2009; Mayo & Scott, 2011), three ⁵³ studies conducted in the United States (Claiborne, 2006; Reeves et al., 2019; Saal et al., 2015), and one study performed in Thailand (Kummarg et al., 2018). The described interventionists who administered the case management interventions in the selected studies included a care coordinator (n = 1 study); nurse case managers ³⁰ 5 studies; a nurse and a physiotherapist (n = 1 study); and an interprofessional team of home care service providers, including a

³⁰ nurse, physiotherapist, occupational therapist, and speech-language pathologist (n = 1 study). The intervention was administered in a hospital (n = 7 studies) and a central community care setting (n = 1 study).

All eight investigations included a total of 1196 stroke survivors. Their ages ranged from 60 to 70.6 years. All studies (n = 8) separated the participants into two groups: those who received normal treatment and those who received the intervention. The intervention group was given a care coordination model that included biopsychosocial concerns, service needs, and self-care practice adherence, as well as evaluations, planning, facilitation, and advocacy for alternatives ⁴¹ resources necessary to fulfill a patient's health requirements. The duration of the intervention ranged from 24 hours to 12 months. The duration of follow-up after intervention ranged from immediately following the intervention to 12 months. Table 1 contains a summary of the included research.

Case management intervention effects for stroke survivors

Mental health

Six studies, which included 930 participants, used the Short Form (SF-36) Health Survey (SF-36) (Claiborne, 2006; Markle-Reid et al., 2011; Mayo et al., 2008), the Mental Health Index (MHI) (Mayo & Scott, 2011), the Patient-Reported Outcomes Measurement Information System-10 (PROMIS-10) (Reeves et al., 2019), and the World Health Organization Quality of Life-BREF (WHOQOL-BREF) (Saal et al., 2015) to measure mental health. The pooled SMD between groups was 0.26 (95% CI: 0.07 to 0.45, $p = 0.01$, $\tau^2 = 0.02$, $I^2 = 43.02\%$; Figure 2; 2.1), indicating that the case management intervention group outperformed the control group in terms of mental health outcomes. Egger's test yielded no significant results ($t = 2.65$, $p = 0.081$).

Activities of daily living

Three studies, which included 555 participants, used the Barthel Index (BI) (Mayo et al., 2008; Mayo et al., 2009) and Instrumental Activities of Daily Living (IADL) (Reeves et al., 2019) to measure the ability to perform activities of daily living. The pooled SMD between groups was 0.68 (95% CI: 0.37 to 0.99, $p < 0.001$, $\tau^2 = 0.05$, $I^2 = 68.61\%$; Figure 2; 2.2), indicating that the case management intervention group outperformed the control group in the performance of activities of daily living. Egger's test resulted in a non-significant result ($t = 0.43$, $p = 0.665$).

Physical function

Three studies, which included 930 participants, SF-36 (Claiborne, 2006; Markle-Reid et al., 2011; Mayo et al., 2009; Mayo & Scott, 2011), the PROMIS-10 (Reeves et al., 2019), and the WHOQOL-BREF (Saal et al., 2015) to measure physical function. The random-effects weighted SMD was 0.05 (95% CI: -0.08 to 0.19, $\tau^2 = 0.00$, $I^2 = 0.00\%$; Figure 2; 2.3), indicating no significant differences in physical function between the intervention and control groups ($p = 0.44$). Egger's test yielded no significant results ($t = 0.76$, $p = 0.448$).

Social function

In three investigations, which included 537 individuals, no significant differences were observed between the intervention and control groups when the SF-36 (Markle-Reid et al., 2011; Mayo & Scott, 2011; Saal et al., 2015) was used to monitor social function. The pooled SMD between groups was 0.33 (95% CI: -0.02 to 0.68, $p = 0.06$, $\tau^2 = 0.07$, $I^2 = 69.78\%$; Figure 2; 2.4). Egger's test yielded no significant results ($t = 0.83$, $p = 0.406$).

Quality assessment for methodology

The RoB-2 was used to evaluate the quality of evidence of the included studies, which were all found to have a low likelihood of bias. The identified high risk of bias domains was primarily linked to the randomization procedure due to a lack of concealment for the intervention groups and the lack of blinding for participants and the intervention provider during participant recruitment. The quality evaluation findings are summarized in Appendix S1. Further, Egger's regression test revealed that the effects of publication bias were minor ($p > 0.05$) for all studies.

DISCUSSION

The results of this meta-analysis suggested that case management interventions substantially outperformed usual care in terms of improving stroke survivors' activities of daily living performance and mental health for 6 weeks to 6 months following the intervention. However, no significant differences in physical or social function were found across groups. Due to the small number of studies that have been conducted within the past decade, the outcomes associated with case management interventions remain equivocal. The present study provides additional evidence that case management may assist stroke survivors throughout the transitional care period. Furthermore, previous studies have shown that case management intervention offers continuity of care when treatment sessions end or follow-up doctor visits become less regular and it appears to benefit a range of different populations than stroke patients (i.e., people with dementia and patients with cardiovascular diseases) (Huntley et al., 2016; Saragih, Tonapa, et al., 2021).

Mental health

Mental health is becoming an increasingly important issue and was examined in six of the eight studies included in this review. Each study used a different case management model, managed by various types of individuals, such as a nurse case manager, a social worker, or as a collaboration between two or more disciplines, and lasting from 6 weeks to 1 year (Claiborne, 2006; Markle-Reid et al., 2011; Reeves et al., 2019; Saal et al., 2015). These studies indicated that stroke survivors who received transitional care via home visits and phone calls had better mental health outcomes than the control group who received standard care. Another study that examined the combination of home visits and mailed letters also showed positive mental health outcomes, including reduced stress and depression levels that lasted for 6 months post-discharge (Ostwald et al., 2014). Stroke survivors also experienced long-term emotional well-being after receiving consultations at a health center six months after stroke (Verberne et al., 2020). This evidence suggests that the mental health of stroke survivors can be supported through various approaches and that these support strategies can be applied at various

43
TABLE 1 Summary of included studies

No	Author, year/country	Health provider	Setting	Participants (experimental/control)		
				Sample size (N)	Female	Age (years old)
1	Clairborne, 2006/USA	Care coordinator	Hospital	28 $N_i = 16$ $N_c = 12$	NA	≥65
2	Kummarg et al., 2018/Thailand	Nurse case manager	Hospital	76 $N_i = 38$ $N_c = 38$	22	≥60
3	Markle-Reid et al., 2011/Canada	Nurse, physiotherapist, occupational therapist, and speech language pathologist	Central Community Care	82 $N_i = 43$ $N_c = 39$	37	70.6
4	Mayo et al., 2008/Canada	Nurse case-manager	Hospital	190 $N_i = 96$ $N_c = 94$	74	70
5	Mayo & Scott, 2011/Canada	Nurse case manager	Hospital	190 $N_i = 96$ $N_c = 94$	48	70
6	Mayo et al., 2009/Canada	Nurse case manager	Hospital	190 $N_i = 96$ $N_c = 94$	NA	70
7	Reeves et al., 2019/USA	Nurse case manager	Hospital	175 $N_i = 88$ $N_c = 87$	48	66
8	Saal et al., 2015/USA	A nurse and a physiotherapist	Hospital	265 $N_i = 130$ $N_c = 135$	128	≥68.1

Abbreviations: BI, Barthel Index; c, control; CNS, Canadian Neurological Scale; GDS, Geriatric Depression Scale; HRQOL, Health-Related Quality of Life; i, intervention; IADL, Instrumental Activities of Daily Living; MHI, Mental Health Index; NA, Not Available; NIHSS, The National Institute of Health Stroke Scale; PBSI, Preference-Based Stroke Index; PHQ, Patient Health Questionnaire; PROMIS-10, Patient-Reported Outcomes Measurement Information System-10; SF-36, Short-Form 36; WHOQOL-BREF, World Health Organization Quality of Life.

times following a stroke to improve mental health outcomes, indicating that any mental health support provided after a stroke could potentially be beneficial.

Activities of daily living

Three studies from the last decade demonstrated that case management interventions managed by individual case managers lasting from 3 weeks to 3 months could effectively improve activities of daily living capacity (Mayo et al., 2008; Mayo et al., 2009; Reeves et al., 2019). Two of these three studies were coordinated by stroke nurses, and the third was managed by a social worker. These studies suggest that stroke nurse specialists may play unique roles in leading and providing individualized stroke-specific transitional care (Chen, Xiao, et al., 2021). Person-centered discharge planning and transitional care managed by stroke nurses have

been suggested to reduce unnecessary hospitalizations for stroke patients during the 30 days following discharge to home (Condon et al., 2016; White et al., 2015). Case management conducted by social workers also resulted in improvements in activities of daily living capacity (Reeves et al., 2019). Both interventions used similar approaches, including home visits and monitoring via telephone calls. The current study found that at the first home visit by neurologic nurses, approximately 64.2% to 91.4% of patients were able to perform activities of daily living without assistance (Pugh et al., 2021). The key treatments for stroke survivors and caregivers following discharge to the home included home visits paired with weekly telephone follow-up and biweekly at-home visits (Lin, Xiao, Chamberlain, Newman, et al., 2020). These interventions allow case managers to continue encouraging participants to track their activities of daily living and progress toward transitional care goals (Lin, Xiao, Chamberlain, Newman, et al., 2020).

Intervention types		Intervention length/follow-up length	Outcomes
Experimental group	Control group		
Received care coordination model; biopsychosocial issues, service needs, and adherence to self-care practices	Usual care	3 months/ Baseline and 3months	Physical and mental component (SF-36); depression (GDS); adherence to self-management practices
Nurse case management	Standard care	24 h/Baseline and 24 h after treatment	Stroke severity (NIHSS)
Received self-management program, promoted behavioral change, and home visit	Usual home care services	12-month/Baseline and 12 months	Cognitive status (SPMSQ); depression (CES-D); Quality of life: physical, mental, social function (SF-36); reintegration normal living (RNLI)
Received the case-management intervention; assessment, planning, facilitation, and advocacy for options and services to meet an individual's health needs through communication and available resources to promote quality cost-effective outcomes	Usual care	6 weeks/6 weeks, and 6 months	Physical and mental component (SF-36); depression (GDS); activities of daily living (BI), reintegration normal living (RNLI); Stroke severity (PBSI)
Received the case-management intervention	Usual care	6 weeks/6 weeks post-intervention, and 6 months	Mental Health (MHI); Physical and social function (SF-36)
Received the stroke Case Manager	Usual Care	6 weeks/6 weeks post-intervention and 6 months	Cognitive status (CNS); Physical and mental component (SF-36), activities of daily living (BI); depression (GDS);
Received self-management program, care coordination, information access, and patient engagement	Usual Care	3 months/1 week and 3 months	Physical and mental health (PROMIS-10); depression (PHQ); Activities of Daily Living (IADL)
Received standard care assessment, and health education; stroke warning signs and symptoms, risk factors and risk management, ambient-assisted living, coping strategies, outpatient therapies, self-help after stroke, and advance care planning	Usual Care	3 months/Baseline and 12 months	Physical, mental health, and social function (WHOQOL-BREF); depression (GDS)

Physical function


The results of the present study demonstrated that case management intervention did not significantly improve physical function (Claiborne, 2006; Markle-Reid et al., 2011; Mayo et al., 2009; Mayo & Scott, 2011; Reeves et al., 2019; Saal et al., 2015). Individual and interprofessional teams conducting home visits and telephone calls are inadequate to achieve desired outcomes. Only one case management intervention applied rehabilitation therapy sessions during the home visit (Saal et al., 2015). A recent comprehensive analysis found that home-based rehabilitation applied in lower-income nations resulted in greater increases in physical function than rehabilitation applied in more industrialized countries (Chi et al., 2020). This phenomenon was influenced by many barriers, ranging from stroke rehabilitation infrastructure and a shortage of physiotherapists and occupational therapists (Pandian et al., 2017). In more developed countries, neurorehabilitation facilities are often equipped with modern, high-technology equipment, such as virtual reality-assisted rehabilitation (Zhang et al., 2021). All of the studies that examined physical function outcomes in this study

were performed in developed countries, such as the United States, which may indicate that stroke survivors in developed countries have access to adequate health care facilities and staff rather than relying on home care (Claiborne, 2006; Reeves et al., 2019; Saal et al., 2015).


Social function

Three studies included in this study assessed the effects of case management treatments on social functioning; however, none of these studies found significant results (Markle-Reid et al., 2011; Mayo & Scott, 2011; Saal et al., 2015). Although not significant, one case management showed clinical improvement among stroke survivors under interprofessional rehabilitation care, which included health care specialists and specialists with non-healthcare backgrounds, such as social workers and personal support workers (Markle-Reid et al., 2011). More research and further developments in the design of interprofessional stroke rehabilitation approaches appear to be promising strategies for improving the


2.1 Mental Health




2.2 Activities of daily living



2.3 Physical function



2.4 Social Function



37

FIGURE 2 Forest plot of case management approaches for stroke survivors

social functions of stroke survivors. Norrving et al. (2018) formed an action plan for supporting the assisted release of stroke survivors in Europe for 2018–2030, which included social and support

workers, in addition to specialized therapists, aiming to provide specialist rehabilitation and social assistance in a home setting (Norrving et al., 2018). This approach appears to be a reasonable

strategy to accommodate the unmet social needs of stroke survivors (McKevitt et al., 2011).

LIMITATIONS

Our study has certain limitations and merits that should be acknowledged. Overall, this study identified that case management might benefit patients' mental health and capacity to perform activities of daily living after stroke. However, more studies remain necessary to establish robust evidence to support case management interventions among stroke survivors because only eight studies were identified during the period from 2006 to 2019, which is an inadequate level of research to validate the current findings. Future trials examining different cultural approaches in different countries remain necessary as stroke remains a global burden. Furthermore, this analysis only included studies published in English; therefore, major findings from studies reported in other languages were excluded. Furthermore, the features of each included study differed significantly. As a result of the small sample size in this study, several statistical analyses, including meta-regression and moderator analyses, for example, could not be performed.

CONCLUSIONS AND RELEVANCE FOR CLINICAL PRACTICE

Finally, this systematic review and meta-analysis found that case management may be useful for improving mental health and the capacity for activities of daily living among stroke survivors. Overall, the results of this study indicate that case management interventions are beneficial for stroke survivors. Case management interventions were designed to guide care among this demographic and are recommended to assist patients during the hospital-to-home transition. This information may be utilized to guide future research when designing strategies for proper care planning among these groups. Future intervention studies to assess the efficacy of case management interventions among stroke survivors are urgently needed to contribute additional information to the evidence base.

CLINICAL RESOURCES

Stroke Signs and Symptoms | cdc.gov

17 AUTHOR CONTRIBUTIONS

Study conception and design: IDS, FHC. **Data collection:** IDS, ISS. **Data analysis and interpretation:** ISS, IDS. **Drafting of the article:** IDS, ISS, DET, SS. **Critical revision of the article:** All authors.

FUNDING INFORMATION

None.

CONFLICT OF INTEREST

None.

5 DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Ita Daryanti Saragih  <https://orcid.org/0000-0002-7177-5933>
Ice Septriani Saragih  <https://orcid.org/0000-0001-9721-378X>
Dame Elysbeth Tuty Arna Uly Tarihoran  <https://orcid.org/0000-0003-2586-1067>
Sapna Sharma  <https://orcid.org/0000-0003-3997-8025>
Fan-Hao Chou  <https://orcid.org/0000-0002-1756-5054>

REFERENCES

- Alonso, A., Aparicio, F. H. J., Benjamin, E. J., Bittencourt, M. S., Callaway, C. W., Carson, F. A. P., Chamberlain, F. A. M., Kissela, B. M., Knutson, F. K. L., & Lee, C. D. (2021). Heart disease and stroke statistics—2021 update. *Circulation*, 143(1), e00-e00.
- Amir-Behghadami, M., & Janati, A. (2020). Population, intervention, comparison, outcomes and study (PICOS) design as a framework to formulate eligibility criteria in systematic reviews. *Emergency Medicine Journal*, 37(6), 387. <https://doi.org/10.1136/emmered-2020-209567>
- Ariful Islam, M., Rahman, A., Aleem, M. A., & Islam, S. M. S. (2016). Prevalence and associated factors of depression among post-stroke patients in Bangladesh. *International Journal of Mental Health and Addiction*, 14(2), 154–166. <https://doi.org/10.1007/s11469-015-9582-x>
- Brazle, F. C. (2019). What is the impact of case management on medical cost reduction in healthcare companies? *Muma Business Review*, 3, 157–163.
- Broderick, J. P., & Abir, M. (2015). Transitions of Care for Stroke Patients. *Circulation: Cardiovascular Quality and Outcomes*, 8(6-suppl_3), S190–S192. <https://doi.org/10.1161/CIRCOUTCOMES.115.002288>
- Cahan, S., & Gamliel, E. (2011). Cohen's d vs alternative standardized mean group difference measures. *Practical Assessment, Research, and Evaluation*, 16(1), 10.
- Camicia, M., Wang, H., DiVita, M., Mix, J., & Niewczyk, P. (2016). Length of stay at inpatient rehabilitation facility and stroke patient outcomes. *Rehabilitation Nursing*, 41(2), 78–90.
- Campbell, B. C. V., & Khatri, P. (2020). Stroke. *The Lancet*, 396(10244), 129–142. [https://doi.org/10.1016/S0140-6736\(20\)31179-X](https://doi.org/10.1016/S0140-6736(20)31179-X)
- Centers for Disease Control and Prevention. (2020). *Stroke signs and symptoms*. National Center for Chronic Disease Prevention and Health Promotion. https://www.cdc.gov/stroke/signs_symptoms.htm
- Chen, L., Xiao, L. D., & Chamberlain, D. (2020). An integrative review: Challenges and opportunities for stroke survivors and caregivers in hospital to home transition care. *Journal of Advanced Nursing*, 76(9), 2253–2265.
- Chen, L., Xiao, L. D., Chamberlain, D., & Newman, P. (2021). Enablers and barriers in hospital-to-home transitional care for stroke survivors and caregivers: A systematic review. *Journal of Clinical Nursing*, 30, 2786–2807.
- Chen, Y., Wei, Y., Lang, H., Xiao, T., Hua, Y., Li, L., Wang, J., Guo, H., & Ni, C. (2021). Effects of a goal-oriented intervention on self-management behaviors and self-perceived burden after acute stroke: A randomized controlled trial. *Frontiers in Neurology*, 12(1166). <https://doi.org/10.3389/fneur.2021.650138>

- Chi, N.-F., Huang, Y.-C., Chiu, H.-Y., Chang, H.-J., & Huang, H.-C. (2020). Systematic review and meta-analysis of home-based rehabilitation on improving physical function among home-dwelling patients with a stroke. *Archives of Physical Medicine and Rehabilitation*, 101(2), 359–373.
- Claiborne, N. (2006). Effectiveness of a care coordination model for stroke survivors: A randomized study. *Health & Social Work*, 31(2), 87–96.
- Clarke, D. J., & Forster, A. (2015). Improving post-stroke recovery: The role of the multidisciplinary health care team. *Journal of Multidisciplinary Healthcare*, 8, 433–442. <https://doi.org/10.2147/JMDH.S68764>
- Clery, A., Bhalla, A., Bisquera, A., Skolarus, L. E., Marshall, I., McKevitt, C., Rudd, A., Sackley, C., Martin, F. C., Manthorpe, J., & Wang, Y. (2020). Long-term trends in stroke survivors discharged to care homes. *Stroke*, 51(1), 179–185. <https://doi.org/10.1161/STROKEAHA.119.026618>
- Condon, C., Lycan, S., Duncan, P., & Bushnell, C. (2016). Reducing readmissions after stroke with a structured nurse practitioner/registered nurse transitional stroke program. *Stroke*, 47(6), 1599–1604.
- Duncan, P. W., Bushnell, C., Sissine, M., Coleman, S., Lutz, B. J., Johnson, A. M., Radman, M., Pvrú Bettger, J., Zorowitz, R. D., & Stein, J. (2021). Comprehensive stroke care and outcomes: Time for a paradigm shift. *Stroke*, 52(1), 385–393.
- Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *BMJ*, 315(7109), 629–634. <https://doi.org/10.1136/bmj.315.7109.629>
- Espenberger, K. R., Fini, N. A., & Peiris, C. L. (2021). Personal and social factors that influence physical activity levels in community-dwelling stroke survivors: A systematic review of qualitative literature. *Clinical Rehabilitation*, 35(7), 1044–1055. <https://doi.org/10.1177/0269215521993690>
- Fabbri, E., De Maria, M., & Bertolaccini, L. (2017). Case management: An up-to-date review of literature and a proposal of a county utilization. *Annals of translational medicine*, 5(20), 396.
- Grant, J. S., Hunt, C. W., & Steadman, L. (2014). Common caregiver issues and nursing interventions after a stroke. *Stroke*, 45(8), e151–e153. <https://doi.org/10.1161/STROKEAHA.114.005094>
- Guo, J., Wang, J., Sun, W., & Liu, X. (2021). The advances of post-stroke depression: 2021 update. *Journal of Neurology*, 269, 1236–1249. <https://doi.org/10.1007/s00415-021-10597-4>
- Hartford, W., Lear, S., & Nimmon, L. (2019). Stroke survivors' experiences of team support along their recovery continuum. *BMC Health Services Research*, 19(1), 723. <https://doi.org/10.1186/s12913-019-4533-z>
- Higgins, J. P. T., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring inconsistency in meta-analyses. *BMJ (Clinical research ed.)*, 327(7414), 557–560. <https://doi.org/10.1136/bmj.327.7414.557>
- Hudon, C., Chouinard, M.-C., Pluye, P., El Sherif, R., Bush, P. L., Rihoux, B., Poitras, M. E., Lambert, M., Zomahoun, H. T. V., & Légaré, F. (2019). Characteristics of case management in primary care associated with positive outcomes for frequent users of health care: A systematic review. *Annals of Family Medicine*, 17(5), 448–458. <https://doi.org/10.1370/afm.2419>
- Huntley, A. L., Johnson, R., King, A., Morris, R. W., & Purdy, S. (2016). Does case management for patients with heart failure based in the community reduce unplanned hospital admissions? A systematic review and meta-analysis. *BMJ Open*, 6(5), e010933. <https://doi.org/10.1136/bmjopen-2015-010933>
- Joo, J. Y., & Liu, M. F. (2017). Case management effectiveness in reducing hospital use: A systematic review. *International Nursing Review*, 64(2), 296–308. <https://doi.org/10.1111/inr.12335>
- Khazaal, W., Taliani, M., Boutros, C., Abou-Abbas, L., Hosseini, H., Salameh, P., & Sadier, N. S. (2021). Psychological complications at 3 months following stroke: Prevalence and correlates among stroke survivors in Lebanon. *Frontiers in Psychology*, 12(2184). <https://doi.org/10.3389/fpsyg.2021.663267>
- Kummarg, U., Sindhu, S., & Muengtawepongpa, S. (2018). The early outcomes of nurse case Management in Patients with acute ischemic stroke treated with intravenous recombinant tissue plasminogen activator: A prospective randomized controlled trial. *Neurology Research International*, 1-8, 1–8. <https://doi.org/10.1155/2018/1717843>
- Liberati, A., Altman, D. G., Tetzlaff, J., Mulrow, C., Gøtzsche, P. C., Ioannidis, J. P. A., Clarke, M., Devereaux, P. J., Kleijnen, J., & Moher, D. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *PLoS Medicine*, 6(7), e1000100. <https://doi.org/10.1371/journal.pmed.1000100>
- Lin, S., Xiao, L. D., & Chamberlain, D. (2020). A nurse-led health coaching intervention for stroke survivors and their family caregivers in hospital to home transition care in Chongqing, China: A study protocol for a randomized controlled trial. *Trials*, 21(1), 240. <https://doi.org/10.1186/s13063-020-4156-z>
- Lin, S., Xiao, L. D., Chamberlain, D., Newman, P., Xie, S., & Tan, J.-Y. (2020). The effect of transition care interventions incorporating health coaching strategies for stroke survivors: A systematic review and meta-analysis. *Patient Education and Counseling*, 103(10), 2039–2060.
- Lindblom, S., Ytterberg, C., Elf, M., & Flink, M. (2020). Perceptive dialogue for linking stakeholders and units during care transitions—a qualitative study of people with stroke, significant others and healthcare professionals in Sweden. *International Journal of Integrated Care*, 20(1), 11.
- Liu, C. J., Chang, W. P., Araujo de Carvalho, I., Savage, K. E. L., Radford, L. W., & Amuthavalli Thiyagarajan, J. (2017). Effects of physical exercise in older adults with reduced physical capacity: Meta-analysis of resistance exercise and multimodal exercise. *International Journal of Rehabilitation Research*, 40(4), 303–314. <https://doi.org/10.1097/mrr.0000000000000249>
- Lukersmith, S., Millington, M., & Salvador-Carulla, L. (2016). What is case management? A scoping and mapping review. *International Journal of Integrated Care*, 16(4), 2. <https://doi.org/10.5334/ijic.2477>
- Markle-Reid, M., Orridge, C., Weir, R., Browne, G., Gafni, A., Lewis, M., Walsh, M., Levy, C., Daub, S., Roberts, J., & Brien, H. (2011). Interprofessional stroke rehabilitation for stroke survivors using home care. *Canadian Journal of Neurological Sciences*, 38(2), 317–334.
- Mayo, N. E., Nadeau, L., Ahmed, S., White, C., Grad, R., Huang, A., Yaffe, M. J., & Wood-Dauphinee, S. (2008). Bridging the gap: The effectiveness of teaming a stroke coordinator with patient's personal physician on the outcome of stroke. *Age and Ageing*, 37(1), 32–38. <https://doi.org/10.1093/ageing/afm133>
- Mayo, N. E., & Scott, S. (2011). Evaluating a complex intervention with a single outcome may not be a good idea: An example from a randomized trial of stroke case management. *Age and Ageing*, 40(6), 718–724. <https://doi.org/10.1093/ageing/afm133>
- Mayo, N. E., Scott, S. C., & Ahmed, S. (2009). Case management post-stroke did not induce response shift: The value of residuals. *Journal of Clinical Epidemiology*, 62(11), 1148–1156. <https://doi.org/10.1016/j.jclinepi.2009.03.020>
- McKevitt, C., Fudge, N., Redfern, J., Sheldenkar, A., Crichton, S., Rudd, A. R., Forster, A., Young, J., Nazareth, I., Rothwell, P. M., & Silver, L. E. (2011). Self-reported long-term needs after stroke. *Stroke*, 42(5), 1398–1403.
- Mendis, S., Davis, S., & Norrving, B. (2015). Organizational update: The world health organization global status report on noncommunicable diseases 2014; one more landmark step in the combat against stroke and vascular disease. *Stroke*, 46(5), e121–e122.
- Mountain, A., Patrice Lindsay, M., Teasell, R., Salbach, N. M., de Jong, A., Foley, N., Bhogal, S., Bains, N., Bowes, R., Cheung, D., & Cameron, J. I. (2020). Canadian stroke best practice recommendations: Rehabilitation, recovery, and community participation following

- stroke. Part two: Transitions and community participation following stroke. *International Journal of Stroke*, 15(7), 789–806. <https://doi.org/10.1177/1747493019897847>
- Norrving, B., Barrick, J., Davalos, A., Dichgans, M., Cordonnier, C., Guekht, A., Kutluk, K., Mikulik, R., Wardlaw, J., Nabavi, D., & Richard, E. (2018). Action plan for stroke in Europe 2018–2030. *European Stroke Journal*, 3(4), 309–336.
- Ostwald, S. K., Godwin, K. M., Cron, S. G., Kelley, C. P., Hersch, G., & Davis, S. (2014). Home-based psychoeducational and mailed information programs for stroke-caregiving dyads post-discharge: A randomized trial. *Disability and Rehabilitation*, 36(1), 55–62.
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., & Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ*, 372, n71. <https://doi.org/10.1136/bmj.n71>
- Pandian, J. D., William, A. G., Kate, M. P., Norrving, B., Mensah, G. A., Davis, S., Roth, G. A., Thrift, A. G., Kengne, A. P., Yu, C., & Kissela, B. M. (2017). Strategies to improve stroke care services in low-and middle-income countries: A systematic review. *Neuroepidemiology*, 49(1–2), 45–61.
- Pugh, J. D., McCoy, K., Needham, M., Jiang, L., Giles, M., McKinnon, E., & Heine, K. (2021). Evaluation of an Australian neurological nurse-led model of postdischarge care. *Health & Social Care in the Community*, 30(4), e962–e973. <https://doi.org/10.1111/hsc.13498>
- Reeves, M. J., Fritz, M. C., Woodward, A. T., Hughes, A. K., Coursaris, C. K., Swierenga, S. J., Nasiri, M., & Freddolino, P. P. (2019). Michigan stroke transitions trial a clinical trial to improve stroke transitions. *Circulation-Cardiovascular Quality and Outcomes*, 12(7), e005493. <https://doi.org/10.1161/circoutcomes.119.005493>
- Reeves, M. J., Hughes, A. K., Woodward, A. T., Freddolino, P. P., Coursaris, C. K., Swierenga, S. J., Schwamm, L. H., & Fritz, M. C. (2017). Improving transitions in acute stroke patients discharged to home: The Michigan stroke transitions trial (MISTT) protocol. *BMC Neurology*, 17(1), 115. <https://doi.org/10.1186/s12883-017-0895-1>
- Rodgers, H., & Price, C. (2017). Stroke unit care, inpatient rehabilitation and early supported discharge. *Clinical Medicine*, 17(2), 173–177.
- Saal, S., Becker, C., Lorenz, S., Schubert, M., Kuss, O., Stang, A., Müller, T., Kraft, A., & Behrens, J. (2015). Effect of a stroke support service in Germany: A randomized trial. *Topics in Stroke Rehabilitation*, 22(6), 429–436. <https://doi.org/10.1179/1074935714Z.0000000047>
- Saragih, I. D., Tarihoran, D. E. T. A. U., Batubara, S. O., Tzeng, H.-M., & Lin, C.-J. (2021). Effects of telehealth interventions on performing activities of daily living and maintaining balance in stroke survivors: A systematic review and meta-analysis of randomised controlled studies. *Journal of Clinical Nursing*, 30, 3139–3152. <https://doi.org/10.1111/jocn.16142>
- Saragih, I. D., Tonapa, S. I., Lin, C.-J., & Lee, B.-O. (2021). Effects of case management intervention for people with dementia and their carers: A systematic review and meta-analysis of experimental studies. *International Journal of Nursing Studies*, 121, 104012. <https://doi.org/10.1016/j.ijnurstu.2021.104012>
- Sedgwick, P., & Marston, L. (2013). Meta-analyses: Standardised mean differences. *BMJ: British Medical Journal*, 347, f7257. <https://doi.org/10.1136/bmj.f7257>
- Towfighi, A., Ovbiagele, B., El Husseini, N., Hackett, M. L., Jorge, R. E., Kissela, B. M., Mitchell, P. H., Skolarus, L. E., Whooley, M. A., & Williams, L. S. (2017). Poststroke depression: A scientific statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*, 48(2), e30–e43. <https://doi.org/10.1161/STR.0000000000000113>
- Tse, T., Yusoff, S., Churilov, L., Ma, H., Davis, S., Donnan, G., & Carey, L. (2017). Increased work and social engagement is associated with increased stroke specific quality of life in stroke survivors at 3 months and 12 months post-stroke: A longitudinal study of an Australian stroke cohort. *Topics in Stroke Rehabilitation*, 24, 1–10. <https://doi.org/10.1080/10749357.2017.1318339>
- Verberne, D., Kroese, M., Staals, J., Ponds, R., & Heugten, C. (2020). Nurse-led stroke aftercare addressing long-term psychosocial outcome: A comparison to care-as-usual. *Disability and Rehabilitation*, 44, 1–9. <https://doi.org/10.1080/09638288.2020.1849417>
- Virani, S. S., Alonso, A., Aparicio, H. J., Benjamin, E. J., Bittencourt, M. S., Callaway, C. W., Carson, A. P., Chamberlain, A. M., Cheng, S., Elkind, M. S., & Dellings, F. N. (2021). Heart disease and stroke statistics—2021 update: A report from the American Heart Association. *Circulation*, 143(8), e254–e743.
- Wang, L., Du, J., Cao, W., & Sun, S. (2021). Trends of stroke attributable to high sodium intake at the global, regional, and national levels from 1990 to 2019: A population-based study. *Neurological Research*, 43(6), 474–481.
- White, C. L., Brady, T. L., Saucedo, L. L., Motz, D., Sharp, J., & Birnbaum, L. A. (2015). Towards a better understanding of readmissions after stroke: Partnering with stroke survivors and caregivers. *Journal of Clinical Nursing*, 24(7–8), 1091–1100.
- Wilson, D. B. (2016). Formulas Used by the " Practical Meta-Analysis Effect Size Calculator. Practical meta-analysis.
- Woodward, J., & Rice, E. (2015). Case management. *Nursing Clinics*, 50(1), 109–121.
- Zhang, B., Li, D., Liu, Y., Wang, J., & Xiao, Q. (2021). Virtual reality for limb motor function, balance, gait, cognition and daily function of stroke patients: A systematic review and meta-analysis. *Journal of advanced nursing*, 77, 3255–3273.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

Appendix S1

How to cite this article: Saragih, I. D., Saragih, I. S., Tarihoran, D. E. T., Sharma, S. & Chou, F.-H. (2023). A meta-analysis of studies of the effects of case management intervention for stroke survivors across three countries. *Journal of Nursing Scholarship*, 55, 345–355. <https://doi.org/10.1111/jnu.12822>

A meta-analysis

ORIGINALITY REPORT

20%

SIMILARITY INDEX

16%

INTERNET SOURCES

13%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	pure.hud.ac.uk Internet Source	1%
2	temp.pedermassor.se.ext.levonline.com Internet Source	1%
3	cris.maastrichtuniversity.nl Internet Source	1%
4	experts.umn.edu Internet Source	1%
5	Submitted to Unizin, LLC Student Paper	1%
6	strokengine.ca Internet Source	1%
7	Zhimin Zheng, Chunxia Li, Fen Fu, Hualing Xie, Yuhua Pan, Jinyu Liao, Xiaoying Chen, Ri-hua Xie, Liqun Yue. "Effectiveness of telehealth interventions on depression symptoms after stroke: A systematic review and meta-analysis", Journal of Affective Disorders, 2023 Publication	<1%

8	www.aminer.cn Internet Source	<1 %
9	Santo Imanuel Tonapa, Aini Inayati, Siriluk Sithichoksakulchai, Ita Daryanti Saragih, Ferry Efendi, Fan-Hao Chou. "Outcomes of nurse-led telecoaching intervention for patients with heart failure: A systematic review and meta-analysis of randomised controlled trials", <i>Journal of Clinical Nursing</i> , 2021 Publication	<1 %
10	www.semanticscholar.org Internet Source	<1 %
11	zenodo.org Internet Source	<1 %
12	Submitted to The Hong Kong Polytechnic University Student Paper	<1 %
13	ebrsr.com Internet Source	<1 %
14	spectrum.diabetesjournals.org Internet Source	<1 %
15	Langduo Chen, Lily Dongxia Xiao, Diane Chamberlain, Peter Newman. "Enablers and barriers in hospital-to-home transitional care for stroke survivors and caregivers: A	<1 %

systematic review", Journal of Clinical
Nursing, 2021

Publication

16

medica-musc.researchcommons.org

Internet Source

<1 %

17

journals.lww.com

Internet Source

<1 %

18

www.jssm.org

Internet Source

<1 %

19

peerj.com

Internet Source

<1 %

20

Zou, Ming-Xiang, Rong-Rong Zhou, Wen-Jun Wu, Ning-Jie Zhang, Wen-En Liu, and Xue-Gong Fan. "The use of pleural fluid procalcitonin and C-reactive protein in the diagnosis of parapneumonic pleural effusions: a systemic review and meta-analysis", The American Journal of Emergency Medicine, 2012.

Publication

<1 %

21

docplayer.net

Internet Source

<1 %

22

Md. Sazedur Rahman, Wenbo Peng, Jon Adams, David Sibbritt. "The use of self-management strategies for stroke

<1 %

rehabilitation: a scoping review", Topics in Stroke Rehabilitation, 2022

Publication

23	doaj.org Internet Source	<1 %
24	assets.researchsquare.com Internet Source	<1 %
25	researchnow-admin.flinders.edu.au Internet Source	<1 %
26	Chiu-Shu Fang, Yu-Kang Tu, Shih-Lun Chang, Chia-Chi Kuo, Ching-Ju Fang, Fan-Hao Chou. "Effectiveness of sound and darkness interventions for critically ill patients' sleep quality: A systematic review and component network meta-analysis", Nursing in Critical Care, 2023 Publication	<1 %
27	Submitted to Salve Regina University Student Paper	<1 %
28	Submitted to Trinity College Dublin Student Paper	<1 %
29	www.cambridge.org Internet Source	<1 %
30	Maureen Markle-Reid, Camille Orridge, Robin Weir, Gina Browne et al. "Interprofessional Stroke Rehabilitation for Stroke Survivors	<1 %

Using Home Care", Canadian Journal of
Neurological Sciences / Journal Canadien des
Sciences Neurologiques, 2014

Publication

31

covid19-data.nist.gov

Internet Source

<1 %

32

"World Stroke Congress Abstracts, 2020",
International Journal of Stroke, 2020

Publication

<1 %

33

bmcmmedicine.biomedcentral.com

Internet Source

<1 %

34

bmjopen.bmj.com

Internet Source

<1 %

35

doc.rero.ch

Internet Source

<1 %

36

"14th World Stroke Congress, Singapore, 26-
29 October 2022", International Journal of
Stroke, 2022

Publication

<1 %

37

9dok.org

Internet Source

<1 %

38

Anna Sidorchuk. "Socioeconomic differences
in lung cancer incidence: a systematic review
and meta-analysis", Cancer Causes & Control,
05/2009

Publication

<1 %

39

Simone Fischer, Jakob Linseisen, Inge Kirchberger, Philipp Zickler, Michael Ertl, Markus Naumann, Christine Meisinger. "Association of post-stroke-depression and health-related quality of life three months after the stroke event. Results from the Stroke Cohort Augsburg (SCHANA) study", *Psychology, Health & Medicine*, 2022

Publication

<1 %

40

academic.oup.com

Internet Source

<1 %

41

www.bmj.com

Internet Source

<1 %

42

Bohan Zhang, Dan Li, Yue Liu, Jiani Wang, Qian Xiao. "Virtual reality for limb motor function, balance, gait, cognition and daily function of stroke patients: A systematic review and meta-analysis", *Journal of Advanced Nursing*, 2021

Publication

<1 %

43

Gumhee Baek, Dain Baik, Nayeon Yi. "The effects of of triage applying artificial intelligence on triage in the emergency department: A systematic review of prospective studies", *Research Square Platform LLC*, 2023

Publication

<1 %

44

J.Y. Joo, M.F. Liu. "Case management effectiveness in reducing hospital use: a systematic review", *International Nursing Review*, 2017

Publication

<1 %

45

Renny Wulan Apriliyasari, Ilham Setyo Budi, Maw Pin Tan, Pei-Shan Tsai. "Physical activity and depression in Indonesian adults with stroke: A nationwide survey", *Journal of Nursing Scholarship*, 2022

Publication

<1 %

46

Shuanglan Lin, Lily Dongxia Xiao, Diane Chamberlain, Peter Newman, Shiqi Xie, Jing-Yu Tan. "The effect of transition care interventions incorporating health coaching strategies for stroke survivors: A systematic review and meta-analysis", *Patient Education and Counseling*, 2020

Publication

<1 %

47

Submitted to Xavier University

Student Paper

<1 %

48

boneandjoint.org.uk

Internet Source

<1 %

49

cyberleninka.org

Internet Source

<1 %

50

d-scholarship.pitt.edu

Internet Source

<1 %

51 media.proquest.com Internet Source <1 %

52 munin.uit.no Internet Source <1 %

53 oa.upm.es Internet Source <1 %

54 repub.eur.nl Internet Source <1 %

55 www.pubfacts.com Internet Source <1 %

56 Mathew J. Reeves, Michele C. Fritz, Amanda T. Woodward, Anne K. Hughes et al. "Michigan Stroke Transitions Trial", *Circulation: Cardiovascular Quality and Outcomes*, 2019
Publication <1 %

57 Michael H. Bloch, Angeli Landeros-Weisenberger, Srijan Sen, Philip Dombrowski et al. "Association of the serotonin transporter polymorphism and obsessive-compulsive disorder: Systematic review", *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 2008
Publication <1 %

58 escholarship.mcgill.ca Internet Source <1 %

escholarship.org

59	Internet Source	<1 %
60	gupea-preproduction.ub.gu.se Internet Source	<1 %
61	journals.plos.org Internet Source	<1 %
62	krex.k-state.edu Internet Source	<1 %
63	ksotjournal.kr Internet Source	<1 %
64	onlinelibrary.wiley.com Internet Source	<1 %
65	oss.signavitae.com Internet Source	<1 %
66	pure.manchester.ac.uk Internet Source	<1 %
67	pureadmin.qub.ac.uk Internet Source	<1 %
68	researchmgt.monash.edu Internet Source	<1 %
69	www.ijpsy.com Internet Source	<1 %
70	Brooke Manderson. "Navigation roles support chronically ill older adults through healthcare"	<1 %

transitions: a systematic review of the literature : Navigation roles: review of literature", Health & Social Care in the Community, 03/2012

Publication

71

Cristina de Diego-Alonso, Julia Blasco-Abadía, Almudena Buesa-Estélez, Rafael Giner-Nicolás et al. "Relationship between Participation in Daily Life Activities and Physical Activity in Stroke Survivors: A Protocol for a Systematic Review and Meta-Analysis", Healthcare, 2023

Publication

<1 %

72

Estêvão Scotti Muzzi Marques Leitão. "Influência de polimorfismos de genes funcionais (GAD1 rs1978340, CACNA1C rs100737 e BDNF rs6265) nos neurometabólitos cerebrais em pacientes com Transtorno Afetivo Bipolar e controles", Universidade de São Paulo. Agência de Bibliotecas e Coleções Digitais, 2023

Publication

<1 %

73

Fuad H. Abuadas, Shahnaz M. Ayasrah, Muayyad M. Ahmad, HanaM. Abu-Snieneh, Hayyaf F. Obiedallah, Iman A. Basheti. "Prevalence of depression and its associated factors among stroke survivors in Saudi Arabia", Nursing Open, 2022

Publication

<1 %

74 Huanyu Mou, Mei Sze Wong, Wai Tong Chien. "Effectiveness of dyadic psychoeducational intervention for stroke survivors and family caregivers on functional and psychosocial health: A systematic review and meta-analysis", *International Journal of Nursing Studies*, 2021
Publication

75 Shuanglan Lin, Shiqi Xie, Jianrong Zhou, Qiang Tu, Chunli Wang, Liping Chen. "Stroke survivors', caregivers' and nurse coaches' perspectives on health coaching program towards hospital-to-home transition care: A qualitative descriptive process evaluation", *Journal of Clinical Nursing*, 2022
Publication

76 digitum.um.es
Internet Source

77 diposit.ub.edu
Internet Source

78 files.eric.ed.gov
Internet Source

79 files.magicapp.org
Internet Source

80 www.informingscience.org
Internet Source

81

Bandak, M., N. Jørgensen, A. Juul, I. R. Vogelius, J. Lauritsen, M. G. Kier, M. S. Mortensen, P. Glovinski, and G. Daugaard. "Testosterone deficiency in testicular cancer survivors - a systematic review and meta-analysis", *Andrology*, 2016.

Publication

<1 %

82

Montgomery, Phyllis, Darren Jermyn, Patricia Bailey, Parveen Nangia, Mary Egan, and Sharolyn Mossey. "Community reintegration of stroke survivors: the effect of a community navigation intervention", *Journal of Advanced Nursing*, 2014.

Publication

<1 %

83

Shuanglan Lin, Chunli Wang, Qiumei Wang, Shiqi Xie, Qiang Tu, Hui Zhang, Mingqian Peng, Jianrong Zhou, Julie Redfern. "The experience of stroke survivors and caregivers during hospital-to-home transitional care: A qualitative longitudinal study", *International Journal of Nursing Studies*, 2022

Publication

<1 %

84

James Lewis, Sarah Marsden, Adrian Cherney, Martine Zeuthen et al. "PROTOCOL: Case management interventions seeking to counter radicalisation to violence: A systematic review of tools and approaches", *Campbell Systematic Reviews*, 2023

<1 %

85

Louise Robinson, Marie Poole, Emma McLellan, Richard Lee et al. "Supporting good quality, community-based end-of-life care for people living with dementia: the SEED research programme including feasibility RCT", Programme Grants for Applied Research, 2020

Publication

<1 %

86

Suzanne H S Lo, Janita P C Chau, Kai C Choi, Rebecca Y M Wong, Jackie C Y Kwan, Issac H L Iu. "Health Professional- and Volunteer-partnered Self-management Support (COMBO-KEY) to Promote Self-efficacy and Self-management Behaviors in People with Stroke: A Randomized Controlled Trial", Annals of Behavioral Medicine, 2023

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On