

# Hotspots sequences

*by* Wani D Gunardi

---

**Submission date:** 09-Nov-2022 05:57AM (UTC+0700)

**Submission ID:** 1948610574

**File name:** Hotspots\_sequences.pdf (2.99M)

**Word count:** 8282

**Character count:** 44301

RESEARCH

Open Access

5

# Hotspots sequences of *gyrA*, *gyrB*, *parC*, and *parE* genes encoded for fluoroquinolones resistance from local *Salmonella* Typhi strains in Jakarta



Ignes Nathania<sup>1</sup>, Ita M. Nainggolan<sup>2,3</sup>, Andi Yasmon<sup>4</sup>, Angela Ch. M. Nusatia<sup>5</sup>, Enty Tjoa<sup>1</sup>, Wani D. Gunardi<sup>6</sup> and Lucky H. Moehario<sup>1\*</sup>

## Abstract

**Background:** Infection of *Salmonella enterica* subsp. *enterica* serovar Typhi is the primary etiology of typhoid fever globally and is common in many developing countries, especially those with dense populations and poor environmental sanitation. Antibiotic fluoroquinolones were used for the treatment in the 1980s due to the resistance to the first-line antibiotics. However, many cases of treatment failure of fluoroquinolones in typhoidal patients have been reported from numerous countries in Asia, Europe, Africa, and America. Mutations in quinolone resistance determining regions (QRDR) genes, *gyrA*, *gyrB*, *parC*, and *parE*, are found in fluoroquinolone-resistant *Salmonella* Typhi. Contrast reports came from the *S. Typhi* isolates in Indonesia, mainly Jakarta and the surroundings, obtained from patients with typhoid fever, with good sensitivity to the fluoroquinolones, i.e., nalidixic acid, ciprofloxacin, moxifloxacin, and levofloxacin. The present study, therefore, aimed to identify the hotspot sequences of *gyrA*, *gyrB*, *parC*, and *parE* genes of the local *S. Typhi* strains based on their susceptibility to fluoroquinolones from patients with typhoid fever in Jakarta and its satellite cities.

**Results:** A total of 28 isolates were identified as *S. Typhi*. All isolates were susceptible to nalidixic acid, levofloxacin, and moxifloxacin. Twenty-seven isolates (96.4%) were susceptible to ciprofloxacin, with one isolate (3.6%) being intermediate. The hotspot sequences of *gyrA*, *gyrB*, *parC*, and *parE* genes from all isolates were identical to the fluoroquinolone-sensitive reference sequence *Salmonella enterica* subsp. *enterica* serovar Typhi Ty2 (NCBI GenBank AE014613.1), including the isolate with intermediate susceptibility. The mutation was not found, and amino acid deduced from all hotspots in susceptible and intermediate isolates showed no replacement in all reported codons.

**Conclusions:** This study showed that the local *S. Typhi* strains from Jakarta and surroundings were susceptible to fluoroquinolones (nalidixic acid, ciprofloxacin, levofloxacin, and moxifloxacin), and the hotspot sequences of the *gyrA*, *gyrB*, *parC*, and *parE* genes were all identical to the reference sequence. Thus, the hotspot sequences of the *gyrA*, *gyrB*, *parC*, and *parE* genes seemingly were conserved in Jakarta's local *S. Typhi* strains and could be considered wild type.

\* Correspondence: lucky.moehario@atmajaya.ac.id

<sup>1</sup> Department of Microbiology, School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia, Jalan Pluit Raya No. 2, North Jakarta 14440, Indonesia

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

The phenotypic susceptibility was consistent with the genotypic characteristic without non-synonymous mutations associated with drug resistance.

**Keywords:** *S. Typhi*, Typhoid fever, Fluoroquinolones, Resistant, QRDR, Hotspots

70

## Background

Typhoid fever is a public health problem with high morbidity and mortality rates mostly in developing countries of Africa, South Asia, and Southeast Asia [1, 2]. Infection of *Salmonella enterica* subsp. *enterica* serovar Typhi is the leading cause of the disease. People living in poverty with poor sanitation system, low water quality, and improper food handling promote higher risk of *S. Typhi* infection. It is estimated that around 13% of the world's population lives in extreme poverty, with a poverty rate of 60% in each country [3]. Data from the Global Burden of Disease (GBD) in 2017 showed the number of *S. Typhi* infections about 10 million cases, with the number of deaths of approximately 116,800. These numbers are much higher than paratyphoid fever cases, around 3 million and 19,100 deaths [4], with the case fatality rate of typhoid fever at 10–30% without treatment and falling to 1–4% with proper treatment [2]. GBD report also revealed the Years of Life Lost (YLL) and Years Lived with Disability (YLD) of typhoid fever were around 8,332,000 and 105,500. Although the trend of typhoid fever decreased each year from 1990 to 2010, the global case still reached 13–20 million cases, with mortality number of 145,000–202,000 cases. Areas with the highest issues were South Asia, East Asia, Southeast Asia, and Sub-Saharan Africa [4].

The only way to treat typhoid fever is through antibiotic therapy. The first-line antibiotics were ampicillin, chloramphenicol, and trimethoprim-sulfamethoxazole, where chloramphenicol became the earliest antibiotic introduced in 1948 on the Malay Peninsula [5]. Ampicillin and trimethoprim-sulfamethoxazole replaced chloramphenicol therapy in the 1970s due to chloramphenicol resistance and severe side effects [5, 6]. The first report of *S. Typhi* strain resistant to chloramphenicol came from England in 1950. Later from 1962 to 1967, chloramphenicol resistance *S. Typhi* strains were also reported from India, West Africa, Greece, and Israel [7]. In 1972, MDR (*multidrug resistance*) *S. Typhi* was reported in Mexico [8]. Since the 1980s, the second-line antibiotic, fluoroquinolones, and third-generation cephalosporins have been recommended to treat MDR (*multidrug resistance*) *S. Typhi*. Alternatively, azithromycin can be applied to cure typhoid fever [9].

In recent years, multiple countries have reported the resistance and decrease-susceptibility of Typhi to fluoroquinolones. The early report came from the Middle

East and Central Asia in 2014, followed by Ghana in 2016 [10, 11]. European Committee on Antimicrobial Susceptibility Testing (EUCAST) database from 2015 revealed that 6% of *Salmonella* isolates were resistant to ciprofloxacin. In 2015 as well, the U.S. National Antimicrobial Resistance Monitoring System (NARMS) announced the rising number of *Salmonella* with decreasing susceptibility to ciprofloxacin from <0.5 to 3.5% since 1996 [64]. Hence, the antibiotic-resistant priority list published by the World Health Organization (WHO) in 2017 showed that fluoroquinolone-resistant *Salmonella* was declared in the “high” category, which meant that novel antibiotics were urgently required to be discovered as alternatives to cure typhoid patients [12]. In the meantime, a study by Moehario et al. 2019 in Indonesia showed a contrasting result that 87.5–100% of *S. Typhi* from Jakarta and Tangerang was yet sensitive to first-line antibiotics and fluoroquinolones [13]. Lugito and Cucunawangsih also revealed similar condition in Karawaci, Tangerang, that the resistance rate of *S. Typhi* isolates to ciprofloxacin and levofloxacin was low [14]. As the capital city, Jakarta is the largest city in Indonesia and Southeast Asia, with over 10 million people inhabited. It is surrounded by satellite towns, namely Bogor, Depok, Tangerang, and Bekasi [15]. Many slums can be found around Jakarta, and these circumstances promote *S. Typhi* dissemination and infection. [45]

Researchers have identified several point mutations in *gyrA*, *gyrB*, *parC*, and *parE*, part of the QRDR. Another mechanism that plays a role in fluoroquinolone-resistant properties is plasmid-mediated. Many plasmid-mediated quinolone resistance (PMQR) genes have been identified i.e. *qnrA*, *qnrB*, *qnrS*, *qnrC*, and *qnrD* [9]. A study by Koirala et al. 2012 [16] and Tack et al. 2019 [17] revealed a point mutation in *gyrA*, *gyrB*, and *parC*. According to a study in Italy by Garcia-Fernández et al. 2015, mutations were detected in *gyrA*, *gyrB*, *parC*, and *parE* [18]. Qian et al. 2020 in China also reported some mutations in all of the QRDR regions [19].

The different situation revealed in Jakarta was possibly caused by the varied length of *S. Typhi* genome. Two studies conducted by Thong et al. 1994 and 1996 have revealed different genome fragment patterns and length of isolates from Malaysia and Papua New Guinea using Pulsed-field Gel Electrophoresis (PFGE) and three restriction endonuclease enzymes [20]. Variation in fragment patterns and genome lengths was reported

further by a study by Moehario et al. 2009 using the PFGE method. The *S. Typhi* isolates from several different areas in Indonesia appeared to have distinct genome lengths ranging from 1.495–4.561 kb and were divided into four clusters. They were from Makassar, Jakarta, Jayapura, and one cluster consisted of diverse areas [21]. On the other hand, the average genome length of >4.300 *S. Typhi* at Typhi Pathogenwatch is 4.7 Mbp [22].

The distinct genome length, global resistance pattern, and the widespread use of antibiotics are predicted to be the main factors that induce *S. Typhi*'s resistance to antibiotics including fluoroquinolones. The hotspot sequences of *gyrA*, *gyrB*, *parC*, and *parE* from Indonesia's local *S. Typhi* strains have not been thoroughly studied. At the same time, most *S. Typhi* isolates in Jakarta and the neighboring towns are still susceptible to fluoroquinolones. This study aimed to examine the hotspot region of the genes encoding fluoroquinolone-resistant *gyrA*, *gyrB*, *parC*, and *parE* in the local Jakarta *S. Typhi* strains based on the fluoroquinolones susceptibility.

## Results

### Isolate identification and antibiotic susceptibility profile

A total of 28 *S. Typhi* isolates from Jakarta and its surroundings collected from 2015 to 2021 were retrieved from  $-20^{\circ}\text{C}$  storage. These isolates originally came from patients with typhoid fever and were treated in hospitals across Jakarta and the surroundings. All of the isolates were re-identified and confirmed as *S. Typhi*. Antibiotic susceptibility from 28 isolates was determined following the guideline from CLSI 2011, CLSI 2020, and EUCAST 2022 [23–25]. All isolates (100%) were susceptible to nalidixic acid, levofloxacin, and moxifloxacin. In contrast, susceptibility to ciprofloxacin was 96.4%, and one isolate (3.6%) showed intermediate susceptibility (Table 1). None of the isolates was resistant to all of the tested fluoroquinolones.

### PCR amplification of *gyrA*, *gyrB*, *parC*, and *parE*

The PCR amplification of the hotspot regions in *gyrA*, *gyrB*, *parC*, and *parE* as the common point mutations related to fluoroquinolones resistance showed DNA fragments that migrated at the position expected for a 381 bp, 444 bp, 564 bp, and 688 bp for each gene respectively (Fig. 1, Fig. 2, Fig. 3, Fig. 4).

### DNA sequencing and amino acid analysis

The DNA sequencing results were aligned with the reference sequence *S. Typhi* Ty2 (NCBI GenBank AE014613.1). The results showed that the hotspot *gyrA* sequences from 28 *S. Typhi* isolates were identical to reference sequence (Fig. 5). Similar results were also found in the hotspot of *gyrB*, *parC*, and *parE* sequences

**Table 1** Antibiotic susceptibility profile of *S. Typhi* isolates in Jakarta

Antibiotic	Sensitive (n = 28)	Intermediate (n = 28)	Resistant (n = 28)
Ciprofloxacin (CLSI 2020)	27 (96.4%)	1 (3.6%)	0
Levofloxacin (EUCAST 2022)	28 (100%)	0	0
Nalidixic acid (CLSI 2011)	28 (100%)	0	0
Moxifloxacin (EUCAST 2022)	28 (100%)	0	0



(Fig. 6, Fig. 7, and Fig. 8). Point mutations in these hotspots were not detected in all genes, not even from the isolate with intermediate ciprofloxacin susceptibility. The sequence data generated in this study have been submitted and are available at NCBI GenBank with the accession numbers listed in Additional file 1.

Codons 82, 83, 87, 119, and 133 in *gyrA* have been reported to have point mutations related to fluoroquinolones resistance. However, our results showed that the amino acid obtained from hotspot *gyrA* sequences in these codons were aspartate (Asp), serine (Ser), aspartate (Asp), alanine (Ala), glycine (Gly), respectively, in all 28 *S. Typhi* isolates, which were in accordance to the reference (Fig. 5).

In the amino acid sequences of *gyrB*, codons 426, 435, 464, 465, 466, and 468 have been reported to have point mutations related to fluoroquinolones resistance. Our study showed contrast results that the amino acid obtained from hotspot *gyrB* sequences in these codons were aspartate (Asp), glycine (Gly), serine (Ser), glutamine (Gln), glutamic acid (Glu), and alanine (Ala), respectively in all 28 *S. Typhi* isolates, which were in accordance to the reference (Fig. 6).

Similar results were also seen in *parC*. The most reported codons in *parC* with point mutations related to fluoroquinolones resistance are codon 79, 80, 84, 92, and 106. On the contrary, our results showed that the amino acid in the hotspot *parC* sequences were identical with the reference. They were aspartate (Asp), serine (Ser), glutamic acid (Glu), proline (Pro), and Tryptophan (Trp), respectively in all 28 *S. Typhi* isolates (Fig. 7).


Although mutations in *parE* related to fluoroquinolones resistance were rarely reported, some researchers have found point mutations in codons 364, 420, 434, 444, and 493. On the other hand, our results showed identical amino acids in the hotspot *parE* sequences compared to the reference, which was alanine (Ala), aspartate (Asp), tyrosine (Tyr), isoleucine (Ile), and serine (Ser), respectively, in all 28 *S. Typhi* isolates (Fig. 8).




**Discussion**

Although multiple countries have announced the distribution of fluoroquinolone-resistant *S. Typhi* and the


related QRDR mutations, the local *S. Typhi* strain in Indonesia, especially Jakarta, still has good susceptibility to fluoroquinolones.




**Fig. 3** Electrophoresis of PCR product from *S. Typhi* hotspot *parC*. M = marker 1 kb, marker size from bottom to top (bp): 250, 500, 750, 1,000, 1,500, 2,000, 2,500, 3,000, 4,000, 5,000, 6,000, 8,000, 10,000. Numbers above the gel correspond to the isolate number in Additional file 1. All of the hotspot *parC* amplicons showed the expected fragments, i.e. 564 bp



**Fig. 4** Electrophoresis of PCR product from *S. Typhi* hotspot *parE*. M = marker 1 kb, marker size from bottom to top (bp): 250, 500, 750, 1,000, 1,500, 2,000, 2,500, 3,000, 4,000, 5,000, 6,000, 8,000, 10,000. Numbers above the gel correspond to the isolate number in Additional file 1. All of the hotspot *parE* amplicons showed the expected fragments, i.e. 688 bp



**Fig. 5** The multiple sequence alignment of hotspot *gyrA* from 28 *S. Typhi* isolates. Black boxes indicated amino acids Asp, Ser, Asp, Ala, and Gly in codons 82, 83, 87, 119, and 133, respectively. Amino acid replacements were not detected on the hotspot *gyrA* in 28 *S. Typhi* isolates. The sequence of *gyrA* can be accessed at NCBI GenBank with the accession number ON220744. Complete accession numbers are available in Additional file 1




**Fig. 6** The multiple sequence alignment of hotspot *gyrB* from 28 *S. Typhi* isolates. Black boxes indicated amino acid Asp, Gly, Ser, Gln, Glu, and Ala in codons 426, 435, 464, 465, 466, and 468, respectively. Amino acid replacements were not detected on the hotspot *gyrB* in 28 *S. Typhi* isolates. The sequence of *gyrB* is available at NCBI GenBank with the accession number ON220772. Complete accession numbers are listed in Additional file 1


16

Sequence mutations in the *gyrA* from fluoroquinolone-resistant *S. Typhi* have been reported from regions with and without endemic typhoid fever. Mutations frequently detected in the hotspot sequences of *gyrA* are shown

in Table 2. These mutations commonly occurred and appeared to be related to the resistance to fluoroquinolones. According to the data collected from several studies, some *S. Typhi* isolates from different countries shared



**Fig. 7** The multiple sequence alignment of hotspot *parC* from 28 S. Typhi isolates. Black boxes indicated amino acid Asp, Ser, Glu, Pro, and Trp in codons 79, 80, 84, 92, and 106, respectively. Amino acid replacements were not detected on the hotspot *parC* in 28 S. Typhi isolates. The sequence of *parC* can be accessed at NCBI GenBank with the accession number ON220800. Complete accession numbers are available in Additional file 1



**Fig. 8** The multiple sequence alignment of hotspot *parE* from 28 S. Typhi isolates. Black boxes indicated amino acid Ala, Asp, Tyr, Ile, and Ser in codons 364, 420, 434, 444, and 493, respectively. Amino acid replacements were not detected on the hotspot *parE* in 28 S. Typhi isolates. The sequence of *parE* can be accessed at NCBI GenBank with the accession number ON220828. Complete accession numbers are available in Additional file 1. The image was cropped in order to show all of the hotspot regions in one frame

similar mutations in the hotspot sequences of *gyrA*. The mutation in Asp82Asn was found solely in Italy [18]. The point mutation in Ser83Phe was found in Congo

[17], Italy [18], India [26–28], Nepal [16, 27], Bangladesh [29], and other South Asian countries [27]. Ser83Tyr and Asp87Asn were distributed in similar countries as



**Table 2** Mutations found in the hotspot *gyrA* of *S. Typhi*

Mutation	Susceptibility Profile	Study
<b>Asp82Asn</b>	NAL (R)	[18]
<b>Ser83Phe</b>	OFX (R), CIP (R), NAL (R)	[16] <sup>f</sup>
	CIP (2)	[17]
	CIP (R), NAL (R)	[18]
	NAL (I) CIP (S)	[18]
	CIP (R), NAL (R)	[18] <sup>a,e</sup>
	NAL (I) CIP (S)	[18] <sup>b</sup>
	NAL (R), CIP (S)	[19] <sup>n</sup>
	NAL (R), CIP (I)	[19] <sup>n,a,p</sup>
	CIP (R), NAL (R)	[26]
	OFX (R), (2) (R), NAL (R)	[27] <sup>g</sup>
	CIP (DS), NAL (R)	[28]
	CIP (R), NAL (R)	[28] <sup>j</sup>
	CIP (DS), NAL (R)	[29] <sup>k</sup>
	CIP (R), NAL (R)	[29] <sup>m</sup>
<b>Ser83Tyr</b>	CIP (DS)	[17]
	NAL (I), CIP (S)	[18] <sup>c</sup>
	(73) (R), CIP (I)	[19]
	NAL (R), CIP (S)	[19]
	CIP (R), NAL (60)	[19] <sup>q</sup>
	OFX (R), CIP (R), NAL (R)	[27] <sup>h</sup>
	CIP (DS), NAL (R)	[28]
	CIP (DS), NAL (R)	[29] <sup>k</sup>
	CIP (2) NAL (R)	[29] <sup>l</sup>
<b>Asp87Asn</b>	CIP (R), NAL (R)	[19]
	NAL (R), CIP (I)	[19] <sup>f</sup>
	NAL (R), CIP (S)	[19] <sup>ts</sup>
	CIP (2) NAL (R)	[19] <sup>s</sup>
	CIP (R), NAL (R)	[18] <sup>d,e</sup>
	NAL (I) CIP (S)	[18] <sup>b</sup>
	CIP (R), NAL (R)	[26]
	OFX (R), CIP (R), NAL (R)	[27] <sup>i</sup>
	CIP (R), NAL (R)	[28] <sup>j</sup>
	CIP (DS), NAL (R)	[29] <sup>k</sup>
<b>Asp87Val</b>	OFX (R), CIP (R), NAL (R)	[16] <sup>f</sup>
<b>Asp87Gly</b>	(30) (DS)	[17]
	NAL (R), CIP (I)	[19] <sup>t</sup>
	CIP (R), NAL (R)	[19] <sup>t</sup>
	CIP (R), NAL (R)	[29] <sup>m</sup>
<b>Ala119Glu</b>	(30) (DS)	[17]
<b>Glu133Gly</b>	NAL (R), CIP (S)	[19] <sup>n,r,u</sup>
	NAL (R), CIP (I)	[19] <sup>n,a,p,r,t</sup>
	CIP (R), NAL (R)	[19] <sup>q,t</sup>
<b>Gly133Glu</b>	NAL (DS)	[30] <sup>v</sup>

<sup>a</sup> Double mutations: *gyrA* Ser83Phe and *gyrB* Gly435Ala; *gyrA* Ser83Phe and *gyrB* Gly435Glu; *gyrA* Ser83Phe and *gyrB* Gly435Val

<sup>b</sup> Four mutations in one isolate: *gyrA* Ser83Phe, *gyrA* Asp87Asn, *gyrB* Gly435Glu, and *parC* Ser80Ile

<sup>c</sup> Double mutations in one isolate: *gyrA* Ser83Tyr and *parE* Ser493Phe

**Table 2** (continued)

<sup>d</sup> Double mutations in one isolate: *gyrA* Asp87Asn and *gyrB* Gly435Glu

<sup>e</sup> Three mutations in one isolate: *gyrA* Ser83Phe, *gyrA* Asp87Asn, and *parC* Ser80Ile

<sup>f</sup> Three mutations: *gyrA* Ser83Phe, *gyrA* Asp87Val, and *parC* Ser80Ile

<sup>g</sup> Three mutations: *gyrA* Ser83Phe, *gyrA* Asp87Val, and *parC* Ser80Ile; Double mutations: *gyrA* Ser83Phe and *parC* Glu84Gly; *gyrA* Ser83Phe and *parE* Ala364Val, in some isolates

<sup>h</sup> Double mutations: *gyrA* Ser83Tyr and *parE* Ala364Val

<sup>i</sup> Double mutations: *gyrA* Asp87Asn and *parE* Ala364Val

<sup>j</sup> Double mutations: *gyrA* Ser83Phe and *gyrA* Asp87Asn

<sup>k</sup> Single mutation with increased efflux pump activity for nalidixic acid

<sup>l</sup> Single mutation with the presence of *qnrS* and increased efflux pump activity for both nalidixic acid and ciprofloxacin

<sup>m</sup> Three mutations: *gyrA* Ser83Phe, *gyrA* Asp87Gly, *parC* Glu92Lys, and increased efflux pump activity for nalidixic acid

<sup>n</sup> Double mutations: *gyrA* Ser83Phe and *gyrA* Glu133Gly

<sup>o</sup> Three mutations: *gyrA* Ser83Phe, *gyrA* Glu133Gly, and *parE* Ile444Ser

<sup>p</sup> Four mutations: *gyrA* Ser83Phe, *gyrA* Glu133Gly, *parE* Ile444Ser, and *parE* Tyr434Ser

<sup>q</sup> Four mutations: *gyrA* Ser83Tyr, *gyrA* Asp87Asn, *gyrA* Glu133Gly, and *parC* Glu84Lys

<sup>r</sup> Double mutations: *gyrA* Asp87Asn and *gyrA* Glu133Gly

<sup>s</sup> Single mutations *gyrA* Asp87Asn and the presence of *aac(6)-Ib-cr4*

<sup>t</sup> Double mutations: *gyrA* Asp87Gly and *gyrA* Glu133Gly

<sup>u</sup> Double mutations: *gyrA* Glu133Gly and *gyrB* Ser426Gly

<sup>v</sup> Double mutations: *gyrA* Ser83Phe and *gyrA* Gly133Glu

Ser83Phe [17, 18, 27–29] with an additional country from China for Asp87Asn [19]. Asp87 had two other mutations, they were Asp87Val, which was found in Nepal [16], and Asp87Gly in Congo [17], China [19], and Bangladesh [29]. Mutations in Ala119Glu, Glu133Gly, and Gly133Glu were only found in Congo [17], China [19], and Iran [30], respectively. Dissimilar identification in the amino acid 133 of *gyrA* was observed in the study by Qian et al. 2020 [19] and Hamidian et al. 2011 [30]. According to the sequence data of *gyrA* *S. Typhi* from NCBI GenBank, the amino acid at position 133 is glycine instead of glutamic acid, as mentioned by Qian et al. 2020 [19]. The amino acid Glu133 was identified in *gyrA* from other *S. enterica* serotypes. Further investigation is needed to confirm the amino acid at position 133. In contrast to the stated reports above, our study showed that all hotspot *gyrA* regions had identical sequences with the reference NCBI GenBank (AE014613.1) at codon Asp82, Ser83, Asp 87, Ala119, and Gly133. Thus, amino acid replacements were not detected in all susceptible and ciprofloxacin intermediate *S. Typhi* isolates.

A similar situation also occurred in *gyrB*, another QRDR gene. Gly435 had three mutations, Gly435Ala, Gly435Glu, and Gly435Val, that were found in Italy. These isolates were from patients with travel histories to Bangladesh and India. Other isolates were isolated from Italian residents that came from Italian, India, and

Bangladesh [18]. In Congo and France, the researcher observed mutation in Ser464Tyr [17, 31]. In the Indian subcontinent, Ser464Phe was reported by Britto et al. 2020 [27]. Despite mutations in amino acids Gln 465Leu, Glu466Asp, and Ala 468Glu, some *S. Typhi* isolates in France showed an interesting profile that was pansusceptible, including fluoroquinolones [31]. Meanwhile, in Congo, Glu466Asp was identified in isolates with decreased susceptibility to ciprofloxacin [17]. Ser426Gly was detected in *S. Typhi* isolates from China only [19]. However, according to the current survey, the amino acid at position 426 was found to be an aspartic acid. Further investigation is needed to confirm the serine amino acid at position 426 of *gyrB*. Although many reports on mutations have been reported earlier, we detected the contrary results from the current study with no point mutations at the hotspot *gyrB* regions, which were Asp426, Gly435, Ser464, Gln465, Glu466, and Ala468. All of them were consistent with the reference sequences. The mutations identified in the hotspot sequences of *gyrB* earlier are summarized in Table 3.

Sequence mutations related to fluoroquinolone-resistant *S. Typhi* in *parC* gene have also been reported, although they were less common than *gyrA* and *gyrB*. Table 4 showed the mutations detected in the hotspot sequences of *parC*.

Only five mutations have been identified in *parC*. A point mutation the amino acid Asp79Gly was found in China from isolates resistant to nalidixic acid but susceptible to ciprofloxacin, as reported by Qian et al. 2020. They also detected mutation in Glu84Lys from isolates resistant to ciprofloxacin and nalidixic acid [19]. In

**Table 3** Mutations found in the hotspot *gyrB* of *S. Typhi*

Mutation	Susceptibility Profile	Study
Ser426Gly	CIP (S), NAL (R)	[19] <sup>1</sup>
Gly435Ala	NAL (R), CIP (R)	[18] <sup>a</sup>
Gly435Glu	NAL (R) CIP (R), NAL (I) CIP (S)	[18] <sup>b</sup>
Gly435Val	NAL (R), CIP (R)	[18] <sup>c</sup>
Ser464Tyr	CIP (DS)	[17]
	NAL (S), CIP (DS)	[31]
Ser464Phe	OFX (R), CIP (R), NAL (R)	[27]
Gln465Leu	Pansusceptible	[31]
Glu466Asp	CIP (DS)	[17]
	Pansusceptible	[31]
Ala468Glu	Pansusceptible	[31]

<sup>1</sup> Double mutations: *gyrA* Glu133Gly and *gyrB* Ser426Gly

<sup>a</sup> Double mutations: *gyrA* Ser83Phe and *gyrB* Gly435Ala

<sup>b</sup> Double mutations: *gyrA* Asp87Asn and *gyrB* Gly435Glu; *gyrA* Ser83Phe and *gyrB* Gly435Glu; Four mutations in one isolate: *gyrA* Ser83Phe, *gyrA* Asp87Asn, *gyrB* Gly435Glu, and *parC* Ser80Ile

<sup>c</sup> Double mutations: *gyrA* Ser83Phe and *gyrB* Gly435Val

**Table 4** Mutations found in the hotspot *parC* of *S. Typhi*

Mutation	Susceptibility Profile	Study
Asp79Gly	NAL (R) CIP (S)	[19]
Ser80Ile	OFX (R), CIP (R), NAL (R)	[16] <sup>1</sup>
Glu84Lys	CIP (R), NAL (R)	[19] <sup>a</sup>
Glu84Gly	OFX (R), CIP (R), NAL (R)	[27] <sup>b</sup>
Glu92Lys	NAL (R) CIP (R), NAL (I) CIP (S)	[29] <sup>c</sup>
Trp106Gly	NAL (R), CIP (R)	[26]

<sup>1</sup> Three mutations: *gyrA* Ser83Phe, *gyrA* Asp87Val, and *parC* Ser80Ile

<sup>a</sup> Four mutations: *gyrA* Ser83Tyr, *gyrA* Asp87Asn, *gyrA* Glu133Gly, and *parC* Glu84Lys

<sup>b</sup> Double mutations: *gyrA* Ser83Phe and *parC* Glu84Gly

<sup>c</sup> Three mutations: *gyrA* Ser83Phe, *gyrA* Asp87Gly, *parC* Glu92Lys, and increased efflux pump activity for nalidixic acid

Nepal, Koirala et al. 2012 identified a mutation in Ser80Ile from isolate resistant to ofloxacin, ciprofloxacin, and nalidixic acid [16]. Chiou et al. 2014 found a mutation in Glu92Lys from an isolate originating from Bangladesh [29]. The glutamic acid at position 92 should be examined since the amino acid proline was considered at this codon. Gopal et al. 2016 showed a mutation in Trp106Gly of *parC* in *S. Typhi* isolate from India resistant to ciprofloxacin and nalidixic acid [26]. The hotspot *parC* regions of all *S. Typhi* isolates from the current study were in line with the reference, in which the codons 79, 80, 84, 92, and 106 were aspartate, serine, glutamic acid, proline, and tryptophan, respectively.

The mutations in *gyrA*, *gyrB*, and *parC* have been mainly reported as the reason for quinolone resistance in *S. Typhi*. On the other hand, only a few reports mentioned the mutations in *parE* regions. The list of reported mutations in hotspot *parE* is shown in Table 5. Accou-Demartin et al. 2011 in France observed mutation at Asp420Asn in *parE* from isolates resistant to nalidixic acid but susceptible to ciprofloxacin [31]. Garcia-Fernández et al. 2015 in Italy reported a mutation in amino acid Ser493Phe in *S. Typhi* isolated from a patient that traveled to India with resistance pattern intermediate susceptibility to nalidixic acid but susceptible to ciprofloxacin [18]. Two mutations in Ile444Ser, and Tyr434Ser of *parE* was found in China by Qian et al. 2020 from *S. Typhi* isolates resistant to nalidixic acid and intermediate susceptibility to ciprofloxacin [19]. In India, Britto et al. 2020 identified one mutation in *parE* Ala364Val from isolates resistant to ciprofloxacin, ofloxacin, and nalidixic acid [27]. Contrastingly, the present study showed amino acids in hotspots *parE* were Ala64, Asp420, Tyr434, Ile444, and Ser493, which were identical to the reference NCBI GenBank (AE014613.1), both in susceptible and intermediate susceptibility isolates to ciprofloxacin.

**Table 5** Mutations found in the hotspot *parE* of *S. Typhi*

Mutation	Susceptibility Profile	Study
Ala364Val	OFX (R), CIP (R), NAL (R)	[27] <sup>a</sup>
Asp420Asn	CIP (S), NAL (R)	[31] <sup>b</sup>
Tyr434Ser	CIP (I), NAL (R)	[19] <sup>c</sup>
Ile444Ser	CIP (I), NAL (R)	[19] <sup>cd</sup>
Ser493Phe	NAL (I), CIP (S)	[18] <sup>e</sup>

<sup>a</sup> Double mutations: *gyrA* Ser83Phe and *parE* Ala364Val; *gyrA* Ser83Tyr and *parE* Ala364Val; *gyrA* Asp87Asn and *parE* Ala364Val

<sup>b</sup> Double mutations: *gyrA* Ser83Phe and *parE* Asp420Asn

<sup>c</sup> Four mutations: *gyrA* Ser83Phe, *gyrA* Glu133Gly, *parE* Ile444Ser, and *parE* Tyr434Ser

<sup>d</sup> Three mutations: *gyrA* Ser83Phe, *gyrA* Glu133Gly, and *parE* Ile444Ser

<sup>e</sup> Double mutations in one isolate: *gyrA* Ser83Tyr and *parE* Ser493Phe

Multiple research across Indonesia supported our results on the susceptibility profile of the local *S. Typhi* strains in Jakarta. *S. Typhi* isolated from 2002 to 2008 in Jakarta and its surroundings showed good susceptibility to ciprofloxacin that was about two decades ago [32] and interestingly a similar profile of susceptibility was found in *S. Typhi* isolated in the late 2010s to 2020s [13, 14]. However, *S. Typhi* isolates collected in Sulawesi, an island located in the East of Indonesia, showed an increasing number of *S. Typhi* resistant to ciprofloxacin [33], and ofloxacin [34].

The distinct strain of *S. Typhi* could be another possible reason for the difference in the susceptibility profile of *S. Typhi* isolates from Jakarta. The study carried out by Chiou et al. 2014 about the genetic relationship of *S. Typhi* from Taiwan, Indonesia, Vietnam, and Bangladesh showed that antibiotic susceptibility is related to clonal variation. Isolates from Bangladesh and Vietnam were closely related with high resistance to nalidixic acid and ciprofloxacin. Isolates from Indonesia were closely related to those from Taiwan and were distant from isolates from Bangladesh and Vietnam. Almost all isolates from Indonesia and Taiwan were sensitive to fluoroquinolones, with a low resistance to nalidixic acid [29]. In 2019, Wang et al. also reported that isolates from Indonesia share the same genotype as the indigenous strain in Taiwan, which was in agreement with the study by Chiou et al. 2014. All isolates in the clade have good susceptibility to ciprofloxacin [35]. Further, the study by Ingle et al. 2019 found three clonalities of *S. Typhi* isolates from Indonesia. One clonality shared the same genotype with isolates from the Philippines, France, and the USA, while the other was also detected in Peru and Bangladesh. All of the *S. Typhi* isolates in these two clonalities were sensitive to fluoroquinolones and had no QRDR mutations, except the isolates from Bangladesh and Peru had a mutation in *gyrA* [36]. Baker et al. 2008 also reported that

no DNA gyrase mutations were detected in *S. Typhi* isolates from Indonesia, although fluoroquinolone-resistant H58 strains have been introduced from neighboring countries [37].

Fluoroquinolones target and convert topoisomerase IV and DNA gyrase. The binding of the antibiotics to both DNA-enzyme complexes thus inhibits the enzyme activity [38]. Mutation in QRDR results in disturbance in replication and separation of DNA, which ultimately will cause cell lysis and cell death [39]. Other mechanisms of fluoroquinolones resistance in bacteria are mediated by plasmids, the so-called PMQR, and chromosome-mediated efflux pump resistance. The expression of PMQR genes has been reported by Qian et al. 2020 through the existence of *aac(6′)-ib-cr-4*, *qnrS1* and *qnrB4* in some *S. Typhi* isolates that were resistant to nalidixic acid and ciprofloxacin [18]. According to Ingle et al. 2019, there were no PMQR genes reported in *S. Typhi* isolates from Indonesia [36]. The existence of PMQR genes in Indonesia has not been reported yet and there are limited reports about the role of efflux pump in fluoroquinolone-resistant *S. Typhi* as well.

## Conclusion

Despite surging cases and mutations in QRDR genes of fluoroquinolone-resistant *S. Typhi* reported worldwide, the overall local strain *S. Typhi* in Jakarta and surroundings showed good susceptibility to fluoroquinolones. Point mutations were not detected in the hotspot *gyrA*, *gyrB*, *parC*, and *parE* areas of all isolates, and no amino acid replacements in all reported codons. All DNA sequences from this study were identical to the reference sequence *S. Typhi* Ty2 (NCBI GenBank AEC4913.1). The absence of point mutations in all hotspots indicated that the phenotypic profile of *S. Typhi* local strains in their susceptibility to ciprofloxacin was consistent with the genotypic characteristic. More research on PMQR genes and efflux pump might be worth to find out the alternative resistance mechanism on intermediate resistance isolate from this research. Further study on resistant local strains *S. Typhi*, if any, would shed more information about mechanisms of resistance to fluoroquinolones.

## Methods

This research was conducted in the Laboratory of Microbiology and Biomolecul, School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia. This study was approved by Ethical Committee of School of Medicine and Health Science, Atma Jaya Catholic University of Indonesia (Ethical approval number: 08/03/KEP-FKIKUAJ/2021 and 9/08/KEP-FKIKUAJ/2021).

### Specimen collection and bacterial identification

In this study, 28 *S. Typhi* isolates from 2015 to 2021 were used. Most of them were recovered from the storage of the Laboratory of Microbiology, School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia. Two isolates were received from hospitals in Jakarta and surroundings. Identification of *S. Typhi* was performed previously using Microbact 12A (Oxoid®) and standard microbiological procedures. Isolates were inoculated in Brain Heart Infusion Broth (BHIB) and Blood Agar (BA). Biochemical test was performed with Triple sugar iron agar (TSIA) and Xylose-lysine-deoxycholate agars (XLD). Positive *S. Typhi* isolate was characterized by the presence of black precipitate and black-centered colony, respectively. Isolates were further identified using Gram staining and serotyping by slide agglutination test with commercial antiserum O9 and antiserum O12 (Difco®). According to the White-Kauffmann Le-Minor scheme classification, *S. Typhi* was identified with positive agglutination. Isolates were then preserved on Nutrient Agar (N76) slant at room temperature. All inoculated media were incubated for 24 h at 35 °C.

### Antibiotic susceptibility test

Antibiotic susceptibility test for ciprofloxacin 5 µg (CIP), nalidixic acid 30 µg (NAL), levofloxacin 5 µg (LVX) and moxifloxacin 5 µg (MXF) was performed with Kirby Bauer or disk diffusion method on Mueller-Hinton agar according to the guideline from Clinical and Laboratory Standards Institute CLSI M100 [23]. The breakpoints of nalidixic acid and ciprofloxacin adhered to the standard from CLSI 2011 and CLSI 2020, respectively [23, 24]. Meanwhile, the breakpoints of levofloxacin and moxifloxacin complied with the data from EUCAST 2022 [25]. *Escherichia coli* ATCC 25922 was used as the reference for the antibiotic susceptibility test.

### DNA extraction

The genomic DNA from all isolates was extracted using QIAmp DNA Mini Kit (Qiagen®) from overnight culture on NA slant according to the instruction from the kit. The extracted DNA was stored in the elution buffer provided by the kit and placed at -20 °C. The presence of the genomic DNA was confirmed by electrophoresis in 0.5% (w/v) agarose gel at 90 V for 60–90 minutes.

### Amplification of *gyrA*, *gyrB*, *parC*, and *parE*

The hotspot regions of *gyrA*, *gyrB*, *parC*, and *parE* listed in Table 2 – Table 5 were amplified by polymerase chain reaction (PCR) using the primers shown in Table 6. These primers were specifically designed for this research according to the reference sequences from NCBI

**Table 6** Primer sequences of *gyrA*, *gyrB*, *parC*, and *parE*

Genes		Size (bp)	Sequences
<i>gyrA</i>	F	381	5'-AAAAATCTGCCGTGTCGTTG-3'
	R		5'-TCACTTCGGTCAGGTGTGTC-3'
<i>gyrB</i>	F	513	5'-AGGTCTGATTGCCGGTGGTTT-3'
	R		5'-AGCTGTCCGGTGTGACTC-3'
<i>parC</i>	F	564	5'-GATCATGGATCGTCCGTGTC-3'
	R		5'-GGCCCTGAACGATATCCAG-3'
<i>parE</i>	F	688	5'-CGCTATGTGCTCTCCGTG-3'
	R		5'-CGCCTTCTCTTCCGTCA-3'

GenBank *S. Typhi* Ty2 (AE014613.1). Hotspot regions were chosen based on the frequent mutations related to fluoroquinolones resistance. PCR was performed using Taq PCR Master Mix Kit (Qiagen®) as specified by the manufacturer's manual with a final reaction volume of 50 µl containing 8 µl (≤1 µg/reaction) of DNA template. The annealing temperature was 60 °C followed by 35x cycles and had been optimized prior to the PCR amplification.

The presence of the targeted fragments was confirmed by electrophoresis, with 4 µl of the PCR product and 1 µl of loading dye loaded in 1.5% (w/v) agarose gel submerged in 1x Tris Boric acid EDTA (TBE). The gel was run at 90 V for 60–90 minutes and stained with Florosafe (1st Base®). The DNA Ladder (Geneaid®) was used to determine the PCR product's molecular size. The DNA fragment was visualized by gel documentation system (Biorad®). Confirmation of gel electrophoresis results was done by 1st Base®.

### DNA sequencing

All PCR products were sequenced by the 1st Base®, Malaysia, using the Sanger sequencing method. SeqTrace and Unipro UGENE were used to analyze the sequences, the results were compared to the reference sequence (NCBI GenBank AE014613.1).

### Abbreviations

S: Sensitive; I: Intermediate; R: Resistant; DC: Decreased susceptibility.

### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12866-022-02666-z>.

**Additional file 1: Table S1.** NCBI Genbank accession numbers and other information related to all hotspot sequences from *S. Typhi* isolates in Jakarta. **Table S2.** Electrophoresis of PCR product from *S. Typhi* hotspot *gyrA* and *gyrB* with the expected fragments of 381 bp and 513 bp, respectively. **Figure S1.** Original electrophoresis image of PCR product from *S. Typhi* hotspot *gyrA* and *gyrB* with the expected fragments of 381

bp and 513 bp, respectively. **Table S3.** Electrophoresis of PCR product from *S. Typhi* hotspot *gyrA* and *gyrB* with the expected fragments of 381 bp and 513 bp, respectively. **Figure S2.** Original electrophoresis image of PCR product from *S. Typhi* hotspot *gyrA* and *gyrB* with the expected fragments of 381 bp and 513 bp, respectively. **Table S4.** Electrophoresis of PCR product from *S. Typhi* hotspot *parC* with the expected fragments of 564 bp. **Figure S3.** Original electrophoresis image of PCR product from *S. Typhi* hotspot *parC* with the expected fragments of 564 bp. **Table S5.** Electrophoresis of PCR product from *S. Typhi* hotspot *parE* with the expected fragments of 688 bp. **Figure S4.** Original electrophoresis image of PCR product from *S. Typhi* hotspot *parE* with the expected fragments of 688 bp. **Table S6.** Electrophoresis of PCR product from *S. Typhi* hotspot *parE* with the expected fragments of 688 bp. **Figure S5.** Original electrophoresis image of PCR product from *S. Typhi* hotspot *parE* with the expected fragments of 688 bp. **Table S7.** Electrophoresis of PCR product from *S. Typhi* hotspot *parE* with the expected fragments of 688 bp. **Figure S6.** Original electrophoresis image of PCR product from *S. Typhi* hotspot *parE* with the expected fragments of 688 bp.

#### Acknowledgements

Not applicable.

#### Authors' contributions

IN: conduct research, primer design, data analysis, manuscript preparation. IMN: primer design, PCR product, sequencing analysis, and review the manuscript. AY: primer design, PCR product and sequencing analysis. ACMN, ET, and WDG: isolate collection. **10** research grand design, determine methods, data analysis, advisory of the research and manuscript. The authors read and approved the final manuscript.

**23**

#### Funding

This work was funded by LPPM of Atma Jaya Catholic University of Indonesia. No. 0139 /III/LPPM-PM.10.01/2/2021 and No. 0487/III/LPPM-PM.10.01/04/2021.

**6**

#### Availability of data and materials

All data generated during this study are included in this published article and available in NCBI Genbank with the links provided in the supplementary information files.

#### Declarations

**25**

#### Ethics approval and consent to participate

This **37** was reviewed and approved by Ethical Committee School of Medicine and Health Sciences of Atma Jaya Catholic University of Indonesia, No. 08/03/KEP-FKIKUA/J/2021 and No. 9/08/KEP-FKIKUA/J/2021.

**14**

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare that they have no conflict of interests.

#### Author details

**4**  
<sup>1</sup>Department of Microbiology, School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia, Jalan Pluit Raya No. 2, North Jakarta 14440, Indonesia. <sup>2</sup>Department of Clinical Pathology, School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia, Jalan Pluit Raya No. 2, North Jakarta 14440, Indonesia. <sup>3</sup>Eijkman Research Center for Molecular Biology, The National Research and Innovation Agency, Jalan Raya Jakarta Bogor No.46, Pakansari, Cibinong, Kabupaten Bogor, Jawa Barat 16915, Indonesia. <sup>4</sup>Department of Microbiology, Faculty of Medicine, Universitas Indonesia, Jalan Pagarangan Timur 16, Jakarta 10320, Indonesia. <sup>5</sup>St. Carolus Hospital, Jalan Sale **33** Raya No. 41, Jakarta 10440, Indonesia. <sup>6</sup>Department of Microbiology, Krida Wacana Christian University, Jalan Arjuna Utara No. 6, Jakarta 11510, Indonesia.

Received: 21 June 2022 Accepted: 10 October 2022

Published online: 18 October 2022

#### References

- Mutai WC, Muigai AWT, Waiyaki P, Kariuki S. Multi-drug resistant *Salmonella enterica* serovar Typhi isolates with reduced susceptibility to ciprofloxacin in Kenya. *BMC Microbiol*. 2018;18:71–5. <https://doi.org/10.1186/s12866-018-1332-3>.
- World Health Organization. Typhoid and other invasive salmonellosis. In: Surveillance standards for vaccine preventable diseases. 2nd ed. Geneva: WHO; 2018.
- Als D, Radhakrishnan A, Arora P, Gaffey MF, Campisi S, Velummailum R, et al. Global trends in typhoidal salmonellosis: a systematic review. *Am J Trop Med Hyg*. 2018;99:10–9. <https://doi.org/10.4269/ajtmh.18-0034>.
- GBD 2017 Typhoid and Paratyphoid Collaborators. The global burden of typhoid and paratyphoid fevers: a systematic analysis for the global burden of disease study 2017. *Lancet Infect Dis*. 2019;19:369–81. [https://doi.org/10.1016/S1473-3099\(18\)30685-6](https://doi.org/10.1016/S1473-3099(18)30685-6).
- Levine MM, Simon R. The Gathering Storm: Is Untreatable Typhoid Fever on the Way? *mBio*. 2018;9(2):e00482–18. <https://doi.org/10.1128/mBio.00482-18>.
- Butler T. Treatment of typhoid fever in the 21st century: promises and shortcomings. *Clin Microbiol Infect*. 2011;17:959–63. <https://doi.org/10.1111/j.1469-0691.2011.03552.x>.
- Anderson ES, Smith HR. Chloramphenicol resistance in the typhoid bacillus. *Br Med J*. 1972;3:329–31. <https://doi.org/10.1136/bmj.3.5822.329>.
- Dyson ZA, Klemm EJ, Palmer S, Dougan G. Antibiotic resistance and typhoid. *Clin Infect Dis*. 2019;68:165–70. <https://doi.org/10.1093/cid/ciy111>.
- Cuyper WL, Jacobs J, Wong V, Klemm EJ, Deborggraeve S, Puyvelde SV. Fluoroquinolone resistance in *Salmonella*: insights by whole-genome sequencing. *Microb Genom*. 2018;7:1–9. <https://doi.org/10.1099/mgen.0.000195>.
- Rahman BA, Wasfy MO, Maksoud MA, Hanna N, Dueger E, House B. Multi-drug resistance and reduced susceptibility to ciprofloxacin among *Salmonella enterica* serovar Typhi isolates from the Middle East and Central Asia. *New Microbes New Infect*. 2014;2:88–92. <https://doi.org/10.1002/nmi.246>.
- Eibach D, Al-Emran HM, Dekker DM, Krumkamp R, Adu-Sarkodie Y, Cruz Espinoza LM, et al. The emergence of reduced ciprofloxacin susceptibility in *Salmonella enterica* causing bloodstream infections in rural Ghana. *Clin Infect Dis*. 2016;62:32–6. <https://doi.org/10.1093/cid/civ757>.
- World Health Organization. WHO publishes list of bacteria for which new antibiotics are urgently needed. 2017. <https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-antibiotics-are-urgently-needed>. Accessed 3 Oct 2021.
- Moehario LH, Robertus T, Tjoa E, Gunardi WD, Nusatia ACM, Edbert D. Antibiotic susceptibility patterns of *Salmonella* Typhi in Jakarta and its trends within the past decade. *J Biol Sci*. 2019;19:40–5. <https://doi.org/10.3923/jbs.2019.40.45>.
- Lugito NPH, Cucunawangsih. Antimicrobial resistance of *Salmonella enterica* serovars Typhi and Paratyphi isolates from a general hospital in Karawaci, Tangerang, Indonesia: a five-year review. *Int J Microbiol*. 2017;1–7. <https://doi.org/10.1155/2017/6215136>.
- World Population Review. Jakarta population 2022. 2022. <https://worldpopulationreview.com/world-cities/jakarta-population>. Accessed 8 June 2022.
- Koirala KD, Thanh DP, Thapa SD, Arjyal A, Karkey A, Dongol S, et al. Highly resistant *Salmonella enterica* Serovar Typhi with a novel *gyrA* mutation raises questions about the long-term efficacy of older fluoroquinolones for treating typhoid fever. *Antimicrob Agents Chemother*. 2012;56:2761–2. <https://doi.org/10.1128/AAC.06414-11>.
- Tack B, Phoba MF, Puyvelde SV, Kalonji LM, Hardy L, Barbe B, et al. *Salmonella* Typhi from blood cultures in the democratic Congo: a 10-year surveillance. *Clin Infect Dis*. 2019;68:130–7. <https://doi.org/10.1093/cid/ciy1116>.
- García-Fernández A, Gallina S, Owczarek S, Dionisi AM, Benedetti I, Decastelli L, et al. Emergence of ciprofloxacin-resistant *Salmonella enterica* serovar Typhi in Italy. *Plos One*. 2015;10:e0132065. <https://doi.org/10.1371/journal.pone.0132065>.
- Qian H, Cheng S, Liu G, Tan Z, Dong C, Bao J, et al. Discovery of seven novel mutations of *gyrB*, *parC* and *parE* in *Salmonella* Typhi and Paratyphi strains from Jiangsu province of China. *Sci Rep*. 2020;10:1–8. <https://doi.org/10.1038/s41598-020-64346-0>.

20. Thong KL, Passey M, Clegg A, Combs BG, Yassin RM, Pang T. Molecular analysis of isolates of *Salmonella* Typhi obtained from patients with fatal and nonfatal typhoid fever. *J Clin Microbiol*. 1996;34:1029–33. <https://doi.org/10.1128/jcm.34.4.1029-1033.1996>.
21. Moehario LH. The molecular epidemiology of *Salmonella* Typhi across Indonesia reveals bacterial migration. *J Infect Dev Ctries*. 2009;3:579–84. <https://doi.org/10.3855/jidc.548>.
22. Argimón S, Yeats CA, Goater RJ, Abudahab K, Taylor B, Underwood A, et al. A global resource for genomic predictions of antimicrobial resistance and surveillance of *Salmonella* Typhi at pathogenwatch. *Nat Commun*. 2021;12(1):2879. <https://doi.org/10.1038/s41467-021-23091-2>.
23. Clinical and Laboratory Standards Institute. CLSI document M100 performance standards for antimicrobial susceptibility testing. 30th ed. Wayne: Clinical and Laboratory Standards Institute; 2020.
24. Clinical and Laboratory Standards Institute. CLSI document M100-S21 performance standards for antimicrobial susceptibility testing twenty-first informational supplement. Wayne: Clinical and Laboratory Standards Institute; 2011.
25. The European Committee on Antimicrobial Susceptibility Testing. Breakpoint tables for interpretation of MIC's and zone diameters version 12.0. 2022. [http://www.eucast.org/clinical\\_breakpoints/](http://www.eucast.org/clinical_breakpoints/). Accessed 8 Aug 2022.
26. Gopal M, Elumalai S, Arumugam S, Durairajpandian V, Kannan MA, Selvam E, et al. GyrA ser83 and parC trp 106 mutations in *Salmonella enterica* serovar Typhi isolated from typhoid fever patients in tertiary care hospital. *J Clin Diagn Res*. 2016;7:14–8. <https://doi.org/10.7860/JCDR/2016/17677.8153>.
27. Britto CD, Dyson ZA, Mathias S, Bosco A, Dougan G, Jose S, et al. Persistent circulation of a fluoroquinolone-resistant *Salmonella enterica* Typhi clone in the Indian subcontinent. *J Antimicrob Chemother*. 2020;75:337–41. <https://doi.org/10.1093/jac/dkz435>.
28. Agrawal P, Kapila K, Kumar S, Ghosh AN, Maurya AK. Novel restriction enzyme S5II for the detection of mutation in *gyrA* gene of *Salmonella enterica* serovar Typhi. *Indian J Pathol Microbiol*. 2010;53:509–12. <https://doi.org/10.4103/0377-4929.68299>.
29. Chiou CS, Lauderdale TL, Phung DC, Watanabe H, Kuo JC, Wang PJ, et al. Antimicrobial resistance in *Salmonella enterica* serovar Typhi isolates from Bangladesh, Indonesia, Taiwan, and Vietnam. *Antimicrob Agents Chemother*. 2014;58:6501–7. <https://doi.org/10.1128/AAC.03608-14>.
30. Hamidian M, Tajbakhsh M, Tohidpour A, Rahbar M, Zali MR, Walther-Rasmussen J. Detection of novel *gyrA* mutations in nalidixic acid-resistant isolates of *Salmonella enterica* from patients with diarrhoea. *Int J Antimicrob Agents*. 2011;37(4):360–4. <https://doi.org/10.1016/j.ijantimicag.2010.12.013>.
31. Accou-Demartin M, Gaborieau V, Song Y, Roumagnac P, Marchou B, Achtman M, et al. *Salmonella enterica* serotype Typhi with nonclassical quinolone resistance phenotype. *Emerg Infect Dis*. 2011;17:1091–4. <https://doi.org/10.3201/eid1706.101242>.
32. Moehario LH, Tjoa E, Kiranasari A, Ningsih I, Rosana Y, Karuniawati A. Trends in antimicrobial susceptibility of gram-negative bacteria isolated from blood in Jakarta from 2002 to 2008. *J Infect Dev Ctries*. 2009;3(11):843–8. <https://doi.org/10.3855/jidc.85>.
33. Hatta M, Ratnawati. Enteric fever in endemic areas of Indonesia: an increasing problem of resistance. *J Infect Dev Ctries*. 2008;2:279–82. <https://doi.org/10.3855/jidc.222>.
34. Jamilah J, Hatta M, Natzir R, Umar F, Sjahril R, Agus R, et al. Analysis of existence of multidrug-resistant H58 gene in *Salmonella enterica* serovar Typhi isolated from typhoid fever patients in Makassar, Indonesia. *New Microbes New Infect*. 2020;38:100793. <https://doi.org/10.1016/j.nmni.2020.100793>.
35. Wang KY, Lee DJ, Shie SS, Chen CJ. Population structure and transmission modes of indigenous typhoid in Taiwan. *BMC Med Genet*. 2019;12(1):126. <https://doi.org/10.1186/s12920-019-0576-6>.
36. Ingle DJ, Nair S, Hartman H, Ashton PM, Dyson ZA, Day M, et al. Informal genomic surveillance of regional distribution of *Salmonella* Typhi genotypes and antimicrobial resistance via returning travellers. *Plos Negl Trop Dis*. 2019;13:e0007620. <https://doi.org/10.1128/AAC.00294-07>.
37. Baker S, Holt K, van de Vosse E, Roumagnac P, Whitehead S, King E, et al. High-throughput genotyping of *Salmonella enterica* serovar Typhi allowing geographical assignment of haplotypes and pathotypes within an urban district of Jakarta, Indonesia. *J Clin Microbiol*. 2008;46(5):1741–6. <https://doi.org/10.1128/JCM.02249-07>.
38. Pham TDM, Ziora ZM, Blaskovich MAT. Quinolone antibiotics. *Med Chem Comm*. 2019;10:1719–39. <https://doi.org/10.1039/c9md00120d>.
39. Aldred KJ, Kerns RJ, Osheroff N. Mechanism of quinolone action and resistance. *Biochemistry*. 2014;53(10):1565–74. <https://doi.org/10.1021/bi5000564>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

### Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)



# Hotspots sequences

---

## ORIGINALITY REPORT

---

18%

SIMILARITY INDEX

12%

INTERNET SOURCES

16%

PUBLICATIONS

5%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- |   |   |     |
|---|---|-----|
| 1 | Axel Dalhoff. "Selective toxicity of antibacterial agents—still a valid concept or do we miss chances and ignore risks?", <i>Infection</i> , 2020<br>Publication  | 2%  |
| 2 | BRENDAN A. NIEMIRA, KELLY A. LONCZYNSKI. "Nalidixic Acid Resistance Influences Sensitivity to Ionizing Radiation among <i>Salmonella</i> Isolates", <i>Journal of Food Protection</i> , 2006<br>Publication | 1%  |
| 3 | <a href="https://journals.plos.org">journals.plos.org</a><br>Internet Source  | 1%  |
| 4 | <a href="http://www.creamjournal.org">www.creamjournal.org</a><br>Internet Source   | 1%  |
| 5 | <a href="http://opac.elte.hu">opac.elte.hu</a><br>Internet Source   | <1% |
| 6 | <a href="http://openaccess.sgul.ac.uk">openaccess.sgul.ac.uk</a><br>Internet Source   | <1% |
| 7 | <a href="http://www.thaiscience.info">www.thaiscience.info</a>  |     |

Internet Source

<1 %

8

[dspace.bracu.ac.bd:8080](https://dspace.bracu.ac.bd:8080)

Internet Source

<1 %

9

Ana Luisa O.C. Carneiro, Rosiane L. Silva, Isabela B.B.E. Rodrigues, Simone C. Machado et al. "Quinolones resistance in Salmonella spp. isolated from broilers and chickens' carcasses under federal inspection", Pesquisa Veterinária Brasileira, 2020

Publication

<1 %

10

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<1 %

11

F Cokro, S T Arrang. "Long-term Proton Pump Inhibitors induces recurrent Urinary Tract Infections: A case study", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

12

C Rodkhum. "gyrA and parC associated with quinolone resistance in Vibrio anguillarum", Journal of Fish Diseases, 5/2008

Publication

<1 %

13

Chiou, C.-S., T.-L. Lauderdale, D. C. Phung, H. Watanabe, J.-C. Kuo, P.-J. Wang, Y.-Y. Liu, S.-Y. Liang, and P.-C. Chen. "Antimicrobial Resistance in Salmonella enterica Serovar Typhi Isolates from Bangladesh, Indonesia,

<1 %



## Taiwan, and Vietnam", Antimicrobial Agents and Chemotherapy, 2014.

Publication

---

14

Jacob Stanley Iramiot, Innocent B. Rwego, Catherine Kansiime, Benon B Asiimwe. "Epidemiology and Antibiotic Susceptibility of *Vibrio cholerae* Associated with the 2017 Outbreak in Kasese district", Research Square Platform LLC, 2019

Publication

---

15

[wprim.whocc.org.cn](http://wprim.whocc.org.cn)

Internet Source

---

16

Arai, K., Y. Hirakata, H. Yano, H. Kanamori, S. Endo, A. Hirotsani, Y. Abe, M. Nagasawa, M. Kitagawa, T. Aoyagi, M. Hatta, M. Yamada, K. Nishimaki, Y. Takayama, N. Yamamoto, H. Kunishima, and M. Kaku. "Emergence of fluoroquinolone-resistant *Streptococcus pyogenes* in Japan by a point mutation leading to a new amino acid substitution", *Journal of Antimicrobial Chemotherapy*, 2011.

Publication

---

17

Lucky Hartati Moehario, Enty Tjoa, Hans Putranata, Shikha Joon, Daniel Edbert, Thomas Robertus. " Performance of TDR-300B and VITEK®2 for the identification of in comparison with VITEK®-MS ", *Journal of International Medical Research*, 2021

<1 %

<1 %

<1 %

<1 %

- 
- |    |   |      |
|----|---|------|
| 18 | Sheetal Shrimal, Anamika Jain. "Synthesis, characterization and biological activity of flavonoid derivatives with amine", AIP Publishing, 2019<br>Publication   | <1 % |
| 19 | <a href="http://aac.asm.org">aac.asm.org</a><br>Internet Source   | <1 % |
| 20 | <a href="http://docsdrive.com">docsdrive.com</a><br>Internet Source   | <1 % |
| 21 | <a href="http://www.science.gov">www.science.gov</a><br>Internet Source   | <1 % |
| 22 | <a href="http://www.cviva.dk">www.cviva.dk</a><br>Internet Source   | <1 % |
| 23 | Daru Seto Bagus Anugrah, Laura Virdy Darmalim, Muhammad Rifky Irwanto Polanen, Permono Adi Putro et al. "Quantum Chemical Calculation for Intermolecular Interactions of Alginate Dimer-Water Molecules", Gels, 2022<br>Publication | <1 % |
| 24 | <a href="http://genomemedicine.biomedcentral.com">genomemedicine.biomedcentral.com</a><br>Internet Source   | <1 % |
| 25 | <a href="http://nagasaki-u.repo.nii.ac.jp">nagasaki-u.repo.nii.ac.jp</a><br>Internet Source   | <1 % |
| 26 | "Abstracts cont.", Clinical Microbiology and Infection, 2004  | <1 % |

27

"Posters", FEMS Microbiology Letters, 2003

Publication

<1 %

---

28

César A. Ribeiro, Dijovani B. dos Reis, Izadora F. Reis, Anderson N. de Carvalho et al.

"Synthesis and antimycobacterial evaluation of fluoroquinolones derivatives coupled with isoprenyl moiety at the C-7 position", Medicinal Chemistry Research, 2022

Publication

<1 %

---

29

C. M. Parry, C. T. Thuy, S. Dongol, A. Karkey et al. "Suitable Disk Antimicrobial Susceptibility Breakpoints Defining Salmonella enterica Serovar Typhi Isolates with Reduced Susceptibility to Fluoroquinolones", Antimicrobial Agents and Chemotherapy, 2010

Publication

<1 %

---

30

M. J. Pons, S. Mosquito, C. Gomes, L. J. del Valle, T. J. Ochoa, J. Ruiz. "Analysis of quinolone-resistance in commensal and diarrheagenic Escherichia coli isolates from infants in Lima, Peru", Transactions of the Royal Society of Tropical Medicine and Hygiene, 2013

Publication

<1 %

---

31

M. Perez-Vazquez, F. Roman, S. Garcia-Cobos, J. Campos. "Fluoroquinolone Resistance in

<1 %

## Haemophilus influenzae Is Associated with Hypermotability", Antimicrobial Agents and Chemotherapy, 2007

Publication

---

32

Mehran Ghaderi, Peter Hjelmström, Göran Hallmans, Fredrik Wiklund, Per Lenner, Joakim Dillner, Carani B Sanjeevi. "MICA gene polymorphism and the risk to develop cervical intraepithelial neoplasia", Human Immunology, 1999

Publication

---

<1 %

33

Submitted to Krida Wacana Christian University

Student Paper

---

<1 %

34

Surojit Das, Sriparna Samajpati, Ujjwayini Ray, Indranil Roy, Shanta Dutta. "Antimicrobial resistance and molecular subtypes of Salmonella enterica serovar Typhi isolates from Kolkata, India over a 15 years period 1998–2012", International Journal of Medical Microbiology, 2017

Publication

---

<1 %

35

Chiou, Chien-Shun, Munirul Alam, Jung-Che Kuo, Yen-Yi Liu, and Pei-Jen Wang. "Chromosome-Mediated Multidrug Resistance in Salmonella enterica Serovar Typhi", Antimicrobial Agents and Chemotherapy, 2015.

Publication

<1 %

---

36 Submitted to Higher Education Commission  
Pakistan  
Student Paper <1 %

---

37 [journal.fk.unpad.ac.id](http://journal.fk.unpad.ac.id)  
Internet Source <1 %

---

38 [www.frontiersin.org](http://www.frontiersin.org)  
Internet Source <1 %

---

39 K. Hirose, A. Hashimoto, K. Tamura, Y.  
Kawamura, T. Ezaki, H. Sagara, H. Watanabe.  
"DNA Sequence Analysis of DNA Gyrase and  
DNA Topoisomerase IV Quinolone Resistance-  
Determining Regions of Salmonella enterica  
Serovar Typhi and Serovar Paratyphi A",  
Antimicrobial Agents and Chemotherapy,  
2002  
Publication <1 %

---

40 [academic.oup.com](http://academic.oup.com)  
Internet Source <1 %

---

41 [core.ac.uk](http://core.ac.uk)  
Internet Source <1 %

---

42 [ddd.uab.cat](http://ddd.uab.cat)  
Internet Source <1 %

---

43 [www.researchgate.net](http://www.researchgate.net)  
Internet Source <1 %

---

44 [www.transpositions.co.uk](http://www.transpositions.co.uk)  
Internet Source <1 %

---

---

45

Ashoka Mahapatra, A. Raj Kumar Patro, Atul Khajuria, Sagarika Dhal, Ashok Kumar Praharaj. "Ciprofloxacin-resistant Gram-negative isolates from a tertiary care hospital in Eastern India with novel gyrA and parC gene mutations", Medical Journal Armed Forces India, 2022

Publication

---

<1 %

46

Guoye Liu, Huimin Qian, Jingwen Lv, Benshun Tian, Changjun Bao, Hong Yan, Bing Gu. "Emergence of mcr-1-Harboring Salmonella enterica Serovar Sinstorf Type ST155 Isolated From Patients With Diarrhea in Jiangsu, China", Frontiers in Microbiology, 2021

Publication

---

<1 %

47

Kumar, Manoj, Sushila Dahiya, Priyanka Sharma, Sujata Sharma, Tej P. Singh, Arti Kapil, and Punit Kaur. "Structure Based In Silico Analysis of Quinolone Resistance in Clinical Isolates of Salmonella Typhi from India", PLoS ONE, 2015.

Publication

---

<1 %

48

Samuel Kariuki, Cecilia Mbae, Robert Onsare, Susan M Kawai et al. " Multidrug-resistant Nontyphoidal Hotspots as Targets for Vaccine Use in Management of Infections in Endemic Settings ", Clinical Infectious Diseases, 2019

Publication

---

<1 %

49

Shao-Chieh Chien, Tsong-Him Iap, Yin-Rong Chiu, Shian-Sen Shie, Chih-Jung Chen.

"Microbiological features of indigenous typhoid cases in Taiwan and relatedness to imported cases, 2001–2014: A cross-sectional analysis", *Travel Medicine and Infectious Disease*, 2018

Publication

<1 %

50

Shinichiro Ota, Yohei Maki, Kazuma Mori, Takaaki Hamamoto et al. "Diagnosis of imported Ugandan typhoid fever based on local outbreak information: A case report", *Journal of Infection and Chemotherapy*, 2016

Publication

<1 %

51

YuQing Liu, JingRan Li, JiaFa Du, Ming Hu, Hua Bai, Jing Qi, Chao Gao, TianTian Wei, Hong Su, JianLing Jin, Peiji Gao. "Accurate assessment of antibiotic susceptibility and screening resistant strains of a bacterial population by linear gradient plate", *Science China Life Sciences*, 2011

Publication

<1 %

52

[microbiomejournal.biomedcentral.com](http://microbiomejournal.biomedcentral.com)

Internet Source

<1 %

53

[oro.open.ac.uk](http://oro.open.ac.uk)

Internet Source

<1 %

54

[pdfs.semanticscholar.org](http://pdfs.semanticscholar.org)

Internet Source

<1 %

---

55

[www.future-science.com](http://www.future-science.com)

Internet Source

<1 %

---

56

"Abstracts cont.", *Clinical Microbiology and Infection*, 4/2005

Publication

<1 %

---

57

"Abstracts of the Asian Congress of Nutrition 2019", *Annals of Nutrition and Metabolism*, 2019

Publication

<1 %

---

58

David J. Edelman, Diah Setyorini Gunawan. "Managing the Urban Environment of Jakarta, Indonesia", *Current Urban Studies*, 2020

Publication

<1 %

---

59

M. A. Pfaller. "Comparison of the broth microdilution (BMD) method of the European Committee on Antimicrobial Susceptibility Testing (EUCAST) with the 24-h CLSI BMD method for fluconazole, posaconazole, and voriconazole susceptibility testing of *Candida* species using epidemiological cutoff values", *Journal of Clinical Microbiology*, 01/12/2011

Publication

<1 %

---

60

M. G. Klotz, Y. C. Kim, J. Katsuwon, A. J. Anderson. "Cloning, characterization and phenotypic expression in *Escherichia coli* of *catF*, which encodes the catalytic subunit of catalase isozyme CatF of *Pseudomonas*

<1 %



# syringae", Applied Microbiology and Biotechnology, 1995

Publication

61

M. MORITA, K. HIROSE, N. TAKAI, J. TERAJIMA et al. " serovar Typhi in Japan, 2001–2006: emergence of high-level fluoroquinolone-resistant strains ", Epidemiology and Infection, 2009

Publication

<1 %

62

Submitted to Udayana University

Student Paper

<1 %

63

Virginia Aiassa, Ana I. Barnes, Andrea M. Smania, Inés Albesa. "Sublethal ciprofloxacin treatment leads to resistance via antioxidant systems in Proteus mirabilis", FEMS Microbiology Letters, 2012

Publication

<1 %

64

[mdpi-res.com](http://mdpi-res.com)

Internet Source

<1 %

65

[mts.intechopen.com](http://mts.intechopen.com)

Internet Source

<1 %

66

[www.freepatentsonline.com](http://www.freepatentsonline.com)

Internet Source

<1 %

67

[www.hindawi.com](http://www.hindawi.com)

Internet Source

<1 %

68

[www.hurights.or.jp](http://www.hurights.or.jp)

Internet Source

<1 %

69

[www.liebertpub.com](http://www.liebertpub.com)

Internet Source

<1 %

70

[www.ndm.ox.ac.uk](http://www.ndm.ox.ac.uk)

Internet Source

<1 %

71

Crump, John A., Maria Sjölund-Karlsson, Melita A. Gordon, and Christopher M. Parry. "Epidemiology, Clinical Presentation, Laboratory Diagnosis, Antimicrobial Resistance, and Antimicrobial Management of Invasive Salmonella Infections", *Clinical Microbiology Reviews*, 2015.

Publication

<1 %

72

*Infectious Diseases and Rural Livelihood in Developing Countries*, 2016.

Publication

<1 %

73

Kaisar A. Talukder, Bijay K. Khajanchi, M. Aminul Islam, Dilip K. Dutta et al. "Genetic relatedness of ciprofloxacin-resistant *Shigella dysenteriae* type 1 strains isolated in south Asia", *Journal of Antimicrobial Chemotherapy*, 2004

Publication

<1 %

74

Lucky H., Enty Tjoa, Veronica N. K. D. Kalay, Angela Abidi. "Chapter 23 Antibiotic

<1 %

## Susceptibility Patterns of Salmonella Typhi in Jakarta and Surrounding Areas", InTech, 2012

Publication

---

75

"Antimicrobial Drug Resistance", Springer Science and Business Media LLC, 2017

Publication

---

<1 %

76

Cecilia S.M. Lucero Estrada, Lidia del Carmen Velázquez, Gabriela Isabel Favier, María Silvia Di Genaro et al. "Detection of Yersinia spp. in meat products by enrichment culture, immunomagnetic separation and nested PCR", Food Microbiology, 2012

Publication

---

<1 %

77

Forcella, C., A. Alessiani, M. Perilli, K. Zilli, E. Di Giannatale, and G. Amicosante. "Characterization of Quinolone Resistance in Escherichia coli Strains of Animal Origin from Italy", Journal of Chemotherapy, 2010.

Publication

---

<1 %

78

J. A. Crump, K. Kretsinger, K. Gay, R. M. Hoekstra et al. "Clinical Response and Outcome of Infection with Salmonella enterica Serotype Typhi with Decreased Susceptibility to Fluoroquinolones: a United States FoodNet Multicenter Retrospective Cohort Study", Antimicrobial Agents and Chemotherapy, 2008

Publication

---

<1 %

79

Kaset, C.. "Rapid identification of lymnaeid snails and their infection with *Fasciola gigantica* in Thailand", *Experimental Parasitology*, 201012

Publication

---

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On